arXiv:nucl-th/9506001v2 12 Mar 1996

A Consistent Meson-Field-Theoretical Description of PP-Bremsstrahlung*

J. A. Eden and M. F. Gari
Institut für Theoretische Physik
Ruhr Universität Bochum, D-44780 Bochum, Germany
email: jamie@deuteron.tp2.ruhr-uni-bochum.de

(31 March 1995, Revised 28 September 1995)

A parameter-free and relativistic extension of the RuhrPot meson-baryon model is used to define the dominant isoscalar meson-exchange currents. We compute pp-bremsstrahlung observables below the $\pi-$production threshold using a relativistic hadronic current density that includes impulse, wave function re-orthonormalization, meson-recoil, $\bar{\mathrm{N}}$N creation and annihilation, $\rho\pi\gamma + \omega\pi\gamma + \rho\eta\gamma + \omega\eta\gamma$ vector-meson decay and N$\Delta\gamma(\pi,\rho)$ exchange currents. We obtain a good description of the available data. The N$\Delta\gamma(\pi)$ current is shown to dominate the large two-body contributions and closed-form expressions for various non-relativistic approximations are analyzed. An experimental sensitivity to the admixture of pseudo-scalar and pseudo-vector admixture of the NNπ interaction is demonstrated. We examine the Lorentz invariance of the NN$\rightleftharpoons$NN t-matrices and show a dominantly pseudo-vector NNπ coupling renders impulse approximation calculations without boost operators to be essentially exact. Conversely, a similar analysis of the ΔN $\rightleftharpoons$NN transitions shows that boost operators and the two-body $N\Delta\gamma$ wave function re-orthonormalization meson-recoil currents are required in NN, ΔN and $\Delta\Delta$ coupled channel t-matrix applications. The need for additional data is stressed.

I. INTRODUCTION

The realization that meson-exchange currents play a vital role in the description of the low-energy pp-bremsstrahlung observables has consequences which are only now coming to be understood. For example, the traditional objective of pp-bremsstrahlung investigations, as indicated in Fig. 1, centers on the capacity of experiment to differentiate the accuracy of the off-shell t-matrices that are predicted by a range of model-dependent NN-interactions. This is now recognized [1,2] as an exceedingly difficult task and, at best, is contingent on a completely reliable and consistent description of the associated meson-exchange currents. As such, a meaningful calculation of the pp-bremsstrahlung observables requires knowledge of the meson-baryon form factors, meson-exchange currents and the NN-interaction within a fully consistent and microscopic effective theory.

Recognizing the importance of exchange currents in pp-bremsstrahlung implies a complete departure form the conventional approach to the problem and considerably changes the nature of such investigation. For many years pp-bremsstrahlung was regarded as something of a special case in nuclear physics because both meson-exchange currents and relativistic effects were expected to be small. The principle reason for this expectation stems from the fact that gauge invariance demands the real photon couples only to conserved currents, so that the n-body parts of the complete hadronic current $J_{[n]}$ for any given NN-interaction V_{NN} must satisfy,



FIG. 1. Relationship between the on-shell t-matrix in pp-scattering, (p,p) and the off-shell t-matrix in pp-bremsstrahlung $(p,p\gamma)$. The parameters in the NN-interaction V defining $t{=}V{+}VGt$ (shown as a bubble) are fitted to the (N,N) phase shifts. Bremsstrahlung is usually considered in nucleon-pole dominance, where (a) initial-, (b) final- and (c) rescattering-interactions are retained, but all meson-exchange currents are neglected. Within this assumption, bremsstrahlung has been regarded as the best means of testing the off-shell t-matrix. However, a sensitivity to off-shell effects requires a large photon energy, so that $G{=}(E-H_0)^{-1}$ diminishes the dominant nucleon pole contributions and the exchange currents become important.

$$
\begin{aligned}
0 &= \nabla.\vec{J} + i\left[\mathcal{H}, J^0\right] \\
\Rightarrow \quad 0 &= \begin{cases} \nabla.\vec{J}_{[1]} + i\left[H_0, J^0_{[1]}\right] & \text{(one body)} \\ \nabla.\vec{J}_{[2]} + i\left[V_{\mathrm{NN}}, J^0_{[1]}\right] & \text{(two body)} \\ +i\left[H_0 + V_{\mathrm{NN}}, J^0_{[2]}\right] & \end{cases}
\end{aligned}
\tag{1.1}
$$

---

*Also to appear in Phys. Rev. **C53**, 1102 (1996)

To isolate the dominant contributions to the observables, it is useful to consider only the static limit, where the two-body charge density $J^0_{[2]}$ can be ignored. The isospin structure of the exchange currents $\vec{J}_{[2]}$ for isovector mesons (π,ρ,..) then reduces to $(\vec{\tau}_1 \times \vec{\tau}_2)^z$, which vanishes in isospin conserving processes like pp-bremsstrahlung. Relativistic processes can also expected to be small since the dominant π-exchange contributions to the N$\bar{\text{N}}$-pair creation and annihilation amplitudes share this isospin structure. Finally, all NNγ couplings with meson-recoil terms can be neglected since they are exactly canceled by the corresponding wave function re-orthonormalization contributions [3]. All of this information suggests that a static limit description of pp-bremsstrahlung involves only the photon coupling to one of the protons either before and/or after (but not during) strong interaction. The leading-order exchange currents, according to this analysis, begin with the η (549 MeV), ω (782 MeV) and ϵ (975 MeV) iso-scalar mesons, and can therefore be reasonably neglected.

However, the above analysis is flawed for several reasons. Even within the static limit there are purely transverse currents $\vec{J}_{\rm t}$ which automatically satisfy $\nabla.\vec{J}_{\rm t}$ =0, so that current conservation places no constraints on the manifestly gauge invariant $\rho\pi\gamma$, $\omega\pi\gamma$ and N$\Delta\gamma$ exchange currents shown in Fig. 2. None of these two-body currents can be included simply by introducing the commutators shown in eq. (1.1), yet they all possess non-vanishing isoscalar contributions which can be important in pp-bremsstrahlung. In addition, as new experiments have succeeded in selecting kinematics that escape the consequences of the low-energy theorems [4,5] and on-shell expansions [6,7], they necessarily emphasize dynamics where the (usually dominant) nucleon-pole contributions of Fig. 1(a-c) are heavily suppressed by the Greens' functions accompanying the highly off-shell t-matrix. As such, otherwise less important contributions gain considerable significance in the observables. This shows that the pp-bremsstrahlung dynamics involves much more than the off-shell t-matrix and the impulse current, and appears to share the complexity of other observables like np-bremsstrahlung and $n + p \rightleftharpoons d + \gamma$.

A very long list of pp-bremsstrahlung calculations have been reported over the last 45 years. We will make no attempt to review them all since more recent works [8–20] already contain appropriate citation and serve to remove a number of questionable approximations. A notable exception to this trend is found in the very detailed r-space calculations reported some 25 years ago by Brown [21,22], where the rescattering amplitudes of Fig. 1(c) were retained and eq. (1.1) was used to constrain the longitudinal meson-exchange currents. Noteworthy calculations since that time have generally been less complete, but find their merit in the application of more reliable NN-interactions and the exploration of coulomb corrections [17] and relativistic corrections to the impulse current [8,11,14,16].



FIG. 2. Currents included in the present calculations: (a) impulse current (b) radiative vector-meson decay currents VPγ = $\rho\pi\gamma$ + $\omega\pi\gamma$ + $\rho\eta\gamma$ + $\omega\eta\gamma$, (c) N$\Delta\gamma$ $\pi-$ and $\rho-$ exchange currents, (d-e) wave function re-orthonormalization and (f) NNγ meson-recoil currents, and (g-i) N$\bar{\text{N}}$-pair creation and annihilation currents. In each of (d-f) and (g-i) we show 3 of the 12 time-ordered diagrams with the energy cuts represented by dotted lines. None of the exchange currents (b-i) can be obtained from eq (1.1).

Only a few of the most recent pp-bremsstrahlung calculations [1,15,19,20] attempt a significant improvement on the standard set by Brown, although there are important technical differences in these works that we will need to consider.

In the present work we will subject the RuhrPot description of meson-baryon interactions to the test of reproducing the pp-bremsstrahlung data below the π-production threshold. The reasons for selecting this effective theory are

- A microscopic definition of the strong form factors is available from non-perturbative and self-consistent calculation [23]. The results are compatible with skyrme [24], non-topological soliton [25] and bag-model [26] calculations, and moreover, with an analysis of experiment [27].

- An NN-interaction model using calculated (not fitted) form factors has been constructed and gives an excellent description of the world data for the NN-scattering phase shifts [28].

- The extension to include meson-exchange currents in the calculation of observables introduces no free parameters whatsoever [1,15].

- A parameter free-extension of the model to define the 3-body interaction has been shown to provide an accurate description of the triton bonding energy [29].

We indicate the relationships between the form factors, the NN interaction and the exchange currents in Fig. 3. Although such consistent calculations can (in principle) be performed for any effective meson field theory, such work has so far only been completed for the RuhrPot description and it appears that there are severe difficulties in obtaining similar consistency in other models. For example, the RuhrPot form factor calculations have been modified to adopt the coupling constants of a conventional boson-exchange NN interaction and yield results [23] which can be accurately parameterized as monopoles with typical regularization scales of $\Lambda \sim 0.8$ GeV. While consistency demands the use of such form factors, conventional boson-exchange models require [30] artificial scales ('cut-offs') of $\Lambda_\pi \geq 1.3$ GeV and $\Lambda_\rho \sim 1.8$ GeV. Such an artificial description of the meson-nucleon vertices necessarily frustrates any attempt to obtain a realistic description of the meson-exchange currents and the 3N interaction.


Electromagnetic Form Factors
$\mathcal{L}_{\mathrm{MB}\gamma}$
Meson-Exchange Currents
Local Gauge Symmetries
Meson-baryon Form Factors
Observables
$\mathcal{L}_{\mathrm{MB}}$
Wave functions
NN interaction

FIG. 3. In the RuhrPot effective meson field theory, meson-baryon form factors are calculated non-perturbatively and the results are used without adjustment as input for subsequent calculations of electromagnetic form factors, the NN interaction and the meson-exchange currents. Such consistency is necessary, for example, to satisfy gauge invariance (which relates the meson-exchange currents to the NN interaction) and ensure orthonormality of the wave functions (through inclusion of wave function re-orthonormalization exchange currents).

Coupled channel t-matrices providing a non-perturbative description of all possible transitions between NN, ΔN and $\Delta\Delta$ states have been available for more than 25 years [31–33] and have already been used to calculate the Δ-isobar contributions to pp-bremsstrahlung observables [19,20]. Under these circumstances it may appear curious that we choose to develop a perturbative description of the N$\Delta\gamma$ $\pi-$ and $\rho-$ exchange currents. However, the coupled channel t-matrices used in recent pp-bremsstrahlung calculations are obtained by the inconsistent combination of the Paris [34] NN$\rightleftharpoons$NN and a static limit version of the Ried-parameterized Bochum [32] NN$\rightleftharpoons$ ΔN interaction. It is therefore impossible to accurately remove the double counted two-pion exchange amplitudes with intermediate NΔ states, so a free parameter is introduced to permit an approximate subtraction procedure. As such, this approach discards from the outset any hope of a obtaining a microscopic description and the quality of the results must be interpreted in terms of a meaningless parameter. We will later show that, even if these inconsistencies were to be resolved by fully consistent calculation, such an approach is contingent upon a reliable description of boost operators, as well as (a subset of) the relativistic meson-exchange currents that will be developed in this work.

In the present work we develop our earlier description [1,14,15] of the RuhrPot meson-baryon interactions in the pp-bremsstrahlung data below the π-production threshold. In refs [14] we included the relativistic single and rescattering impulse-current amplitudes, and in refs [1,15] we introduced the fully relativistic description of the radiative vector-meson decay currents and the non-relativistic description of the N$\Delta\gamma$ $\pi-$ and $\rho-$meson exchange currents without recourse to the soft-photon approximation. In the present work we investigate a number of important extensions. In particular, after describing our model-independent formalism in sec II, we provide in sec III the first bremsstrahlung calculations including a fully relativistic description of the wave function re-orthonormalization and meson-recoil currents that are required to ensure the orthonormality of the wave functions is preserved. We also investigate the Lorentz structure of the NNπ vertex by providing the first calculations for the purely relativistic N$\bar{\mathrm{N}}$ pair creation and annihilation currents. We further present relativistic expressions for the N$\Delta\gamma$ $\pi-$ and $\rho-$exchange currents and identify the source of error in various approximations. Supporting calculation details are supplied in the three appendices. In sec IV, after establishing the sensitivity of the selected pp-bremsstrahlung observables to each of these currents and concluding that a relativistic calculation of the isobar amplitudes is necessary, we compare our relativistic results with the complete data set available from the 1990 TRIUMF pp-bremsstrahlung experiment [35]. We obtain a good description of the experimental data and conclude that a large pseudoscalar admixture in the NNπ Lagrangian is ruled out. Further conclusions and future objectives are given in sec V.

II. FORMALISM

A. Observables

We begin by presenting the model independent expressions we require for the calculation observables for the reaction $N+N \rightarrow N+N+\gamma$. The S–matrix from covariant perturbation theory

$$S_{fi} = \delta_{fi} - i \int d^4x \langle f | J^{\mu}(x) \mathcal{A}_{\mu}(x) | i \rangle + \cdots \qquad (2.1)$$

gives the probability amplitude for a transition $|i\rangle \rightarrow |f\rangle$ as a series involving 0,1,... interactions where the photon field $\mathcal{A}_{\mu}(x)$ couples to the hadronic current density $J^{\mu}(x)$. Since only terms with an odd number of electromagnetic interactions can contribute to the production of a single real photon, and each of these diminishes by $\alpha \sim 1/137$, we retain only the lowest-order contribution and define the transition amplitude as,

$$(2\pi)^4 \mathcal{T}_{fi} \delta^{(4)}(P_f - P_i) = -i \int d^4x \langle f | J^{\mu}(x) \mathcal{A}_{\mu}(x) | i \rangle \qquad (2.2)$$

Following a well trodden path, we integrate the transition amplitude over the phase space available to the final state, and divide by the incident flux, so that with plane wave normalized to a δ-function, we obtain the Lab-frame differential cross section as,

$$\frac{d^3\sigma}{d\Omega_3 d\Omega_4 d\theta_{\gamma}} = \frac{(2\pi)^{-5}\frac{1}{2}m^3}{|\vec{p}_1|} |\, \widetilde{\mathcal{M}}_{fi} \,|^2 J_{\rm ps} \qquad (2.3)$$

where, for the pp- and nn-bremsstrahlung reactions we have

$$|\, \widetilde{\mathcal{M}}_{fi} \,|^2 = \begin{cases} |\mathcal{M}^2_{fi}| & \text{if } M_{S_i},\, M_{S_f},\, \lambda \text{ measured} \\ \dfrac{1}{4} \displaystyle\sum_{S_i M_{S_i}} \sum_{S_f M_{S_f}} \sum_{\lambda=1}^{2} |\mathcal{M}_{fi}|^2 & \text{if } M_{S_i},\, M_{S_f},\, \lambda \text{ not measured} \end{cases} \qquad (2.4)$$

and for np-bremsstrahlung we require,

$$|\, \widetilde{\mathcal{M}}_{fi} \,|^2 = \begin{cases} \dfrac{1}{2} \displaystyle\sum_{T_i, T_f} |\mathcal{M}_{fi}|^2 & \text{if } M_{S_i},\, M_{S_f},\, \lambda \text{ measured} \\ \dfrac{1}{8} \displaystyle\sum_{T_i, T_f} \sum_{S_i M_{S_i}} \sum_{S_f M_{S_f}} \sum_{\lambda=1}^{2} |\mathcal{M}_{fi}|^2 & \text{if } M_{S_i},\, M_{S_f},\, \lambda \text{ not measured} \end{cases} \qquad (2.5)$$

with the invariant amplitude given by,

$$\mathcal{M}_{fi} = i(2\pi)^{15/2} m^{-2} \left[2\omega E_1 E_2 E_3 E_4\right]^{1/2} \mathcal{T}_{fi} \qquad (2.6)$$

We retain only the transverse polarization vectors for the real photon since, within the Gupta-Bleuler quantization formalism, the longitudinal and scalar components can be made to cancel with a gauge transformation, and therefore cannot effect the observables. The 'phase-space factor' $J_{\rm ps}$ appearing in eq (2.3) is defined for arbitrary non-coplanarity $\Phi = (\pi + \phi_3 - \phi_4)/2$ as,

$$J_{\rm ps} = \frac{p_3^2 p_4^2}{E_3 E_4 |\cos\theta_{\gamma}|} \left| \frac{N}{k \sin\theta_{\gamma} \cos\theta_{\gamma}} \right|^{-1} \qquad (2.7)$$

with

$$\begin{aligned} N = {} & \big(p_4 \sin\theta_4 - p_3 \sin\theta_3\big)\big[\sin(\theta_3 + \theta_4) \\ & \quad - \big(\beta_3 \sin\theta_4 + \beta_4 \sin\theta_3\big)\cos\theta_{\gamma}\big] \\ & - k \sin^2\theta_{\gamma} \big(\beta_3 \cos\theta_4 - \beta_4 \cos\theta_3\big) \\ & + 2 \sin\theta_3 \sin\theta_4 \sin^2\Phi \big[\big(p_3 \cos\theta_3 - p_4 \cos\theta_4\big) \\ & \quad - \big(p_3\beta_3 - p_4\beta_4\big)\cos\theta_{\gamma}\big] \end{aligned} \qquad (2.8)$$

where we use $\beta_i = p_i/E_i$ for laboratory-frame reaction kinematics $p_1 + p_2 \rightarrow p_3 + p_4 + k$. We realize that in the limit $\theta_{\gamma} \rightarrow 0$, where $\Phi \rightarrow 0$ is guaranteed, N^{-1} possesses singularities but $k \sin\theta_{\gamma}/N$ remains well defined. Our description of kinematics and phase space is the same as that reported in the detailed discussion of ref [36]. As such, it is sufficient to note that in ref [36] it was shown that $J_{\rm ps}$ possess a square root singularity at the kinematic limit of non-coplanarity, although in the present work we avoid the non-relativistic simplifications discussed therein.

For the calculation of polarized observables it is convenient to denote $d\sigma(\pm\hat{i})$ as the cross section of eq (2.3) measured with the beam polarized in the $\pm\hat{i}$ direction. We choose the quantization axis as the beam direction in the lab frame, and define the vector analyzing powers as,

$$A_i = \frac{d\sigma(+\hat{i}) - d\sigma(-\hat{i})}{d\sigma(+\hat{i}) + d\sigma(-\hat{i})} = \frac{\sum_{T_i T_f \lambda} \mathrm{Tr}\Big\{ \mathcal{M}_{fi} (\vec{\sigma}.\hat{i})_{[1]} \mathcal{M}^{\dagger}_{fi} \Big\}}{\sum_{T_i T_f \lambda} \mathrm{Tr}\Big\{ \mathcal{M}_{fi} \mathcal{M}^{\dagger}_{fi} \Big\}} \qquad (2.9)$$

where $i = \hat{x}, \hat{y}$ or $\hat{z}$ in the lab frame. Similarly, the tensor analyzing powers (sometimes called 'spin-correlation coefficients') are given by,

$$A_{ij} = \frac{d\sigma(+\hat{i},+\hat{j}) + d\sigma(-\hat{i},-\hat{j}) - d\sigma(+\hat{i},-\hat{j}) - d\sigma(-\hat{i},+\hat{j})}{d\sigma(+\hat{i},+\hat{j}) + d\sigma(-\hat{i},-\hat{j}) + d\sigma(+\hat{i},-\hat{j}) + d\sigma(-\hat{i},+\hat{j})}$$

$$= \frac{\sum_{T_iT_f\lambda} \mathrm{Tr}\Big\{\mathcal{M}_{fi}(\vec{\sigma}.\hat{\imath})_{[1]}(\vec{\sigma}.\hat{\jmath})_{[2]}\mathcal{M}^{\dagger}_{fi}\Big\}}{\sum_{T_iT_f\lambda} \mathrm{Tr}\Big\{\mathcal{M}_{fi}\mathcal{M}^{\dagger}_{fi}\Big\}} \tag{2.10}$$

where, for example, $d\sigma(+\hat{\imath}, -\hat{\jmath})$ is the cross section measured with the beam polarized in the $+\hat{\imath}$ direction and the target polarized in the $-\hat{\jmath}$ direction.

B. The Hadronic Current

To obtain a microscopic definition of the invariant amplitude we compute the Fock-space matrix elements of the photon field and hadronic current densities appearing in eq (2.2), so that after making use of translational invariance and selecting the Lorentz-Heaviside system with natural units, eq (2.6) can be recast as,

$$\mathcal{M}_{fi} = (2\pi)^6 m^{-2}\sqrt{E_1E_2E_3E_4}\,\epsilon_\mu(\vec{k},\lambda)\langle\Psi^{(-)}_f|J^\mu(0)|\Psi^{(+)}_i\rangle \tag{2.11}$$

where the field-theoretic hadronic current is given by,

$$J^\mu(x) = \partial_\nu\frac{\partial\mathcal{L}}{\partial(\partial_\nu\mathcal{A}_\mu)} - \frac{\partial\mathcal{L}}{\partial\mathcal{A}_\mu} \tag{2.12}$$

for the Lagrangian $\mathcal{L}$ describing electromagnetic interactions with the interacting meson-baryon system. Direct calculation of eq. (2.11) is impossible since $|\Psi^{(+)}_i\rangle$ and $|\Psi^{(-)}_f\rangle$ represent complete meson-baryon states and therefore involve nucleon, resonance and meson degrees of freedom to infinite order. The problem can, however, be approached with a Hamilton formalism [3] where the total Hilbert space is partioned into meson+resonance+anti-nucleon vacuum- and existing subspaces,

$$\mathcal{H}_\eta = \Big\{|NN\rangle\Big\}, \quad \mathcal{H}_\lambda = \Big\{|\text{the rest}\rangle\Big\} \tag{2.13}$$

We will refer to these subspaces as the η-space and λ-space respectively. Defining η and λ as operators satisfying the conventional algebra $\eta^2 = \eta$, $\lambda^2 = \lambda$ and $\eta\lambda = \lambda\eta = 0$ and which project out the components of $\mathcal{H}_\eta$ and $\mathcal{H}_\lambda$ respectively, we apply a unitary transformation to decouple the meson-resonance vacuum and existing components of wave functions. Although this formalism makes provision for applications involving (for example) explicit meson and/or Δ (see App. A) degrees of freedom in the initial and final states, we confine our present application to energies below the π-production threshold, so that the complete interacting meson-baryon wave function can be written as,

$$|\Psi\rangle = (1 + A)\frac{1}{\sqrt{1 + A^\dagger A}}|\mathcal{X}\rangle \tag{2.14}$$

where $|\mathcal{X}\rangle$ is the two-nucleon state vector and $A = \lambda A\eta$ is required to satisfy $\lambda(H + [H, A] - AHA)\eta = 0$. In particular, we can expand both A and the Hamiltonian H in powers n of the coupling constant, so that with the free-particle energy denoted as H_0, we have,

$$H = H_0 + \sum_{n=1}^{\infty} H_n, \qquad A = \sum_{n=1}^{\infty} A_n \tag{2.15}$$

and note that $\eta H_I\eta{=}\eta H_0\lambda{=}\lambda H_0\eta{=}0$, to obtain,

$$\begin{aligned} 0 = \sum_{n=1}^{\infty}\lambda\Bigg[&H_n + [H_0\,, A_n] + \sum_{i=1}^{n-1} H_iA_{n-i} \\ &- \sum_{i=1}^{n-2}\sum_{j=1}^{n-i-1} A_iH_jA_{n-i-j}\Bigg]\eta \end{aligned} \tag{2.16}$$

We are free to further constrain A by demanding eq. (2.16) is satisfied at each order of n, as would be required for any perturbative application. Since $H_0\,\eta|\Psi\rangle{=}\mathcal{E}_i\,\eta|\Psi\rangle$, where $\mathcal{E}_i$ is the asymptotic energy of the free two-nucleon state, we obtain,

$$\begin{aligned} (\mathcal{E}_i - H_0)A_n = \lambda\Bigg[&H_n + \sum_{i=1}^{n-1} H_iA_{n-i} \\ &- \sum_{i=1}^{n-2}\sum_{j=1}^{n-i-1} A_iH_jA_{n-i-j}\Bigg]\eta \end{aligned} \tag{2.17}$$

Since $A_0{=}0$, we have

$$A_1 = \frac{\lambda}{\mathcal{E}_i - H_0}H_1\eta \tag{2.18a}$$

$$A_2 = \frac{\lambda}{\mathcal{E}_i - H_0}H_2\eta + \frac{\lambda}{\mathcal{E}_i - H_0}H_1\frac{\lambda}{\mathcal{E}_i - H_0}H_1\eta \tag{2.18b}$$

$$\vdots$$

Finally, with

$$H_I = -\int\mathcal{L}d^3x \tag{2.19}$$

we observe that A is completely determined by the strong interaction Lagrangian density defining any model of interest. Combining eqs. (2.11) and (2.14), we then obtain,

$$\mathcal{M}_{fi} = (2\pi)^6 m^{-2}\sqrt{E_1E_2E_3E_4}\langle\mathcal{X}_f|\,\epsilon_\mu(\vec{k},\lambda)J^\mu_{\text{eff}}(0)\,|\mathcal{X}_i\rangle \tag{2.20}$$

where

$$\begin{aligned} J^\mu_{\text{eff}}(0) &= \eta\frac{1}{\sqrt{1 + A^\dagger A}}(1 + A^\dagger)J^\mu(0)(1 + A)\frac{1}{\sqrt{1 + A^\dagger A}}\eta \\ &= \eta\big[J^\mu(0) + J^\mu(0)A + A^\dagger J^\mu(0) + A^\dagger J^\mu(0)A \\ &-\frac{1}{2}J^\mu(0)A^\dagger A - \frac{1}{2}A^\dagger AJ^\mu(0) + \cdots\big]\eta \end{aligned} \tag{2.21}$$

is the effective meson-baryon current density. Eq. (2.21) provides a time-ordered relativistic description of the impulse- and meson-exchange currents implied by the strong interaction Lagrangian density defining any model of interest. This provides, without approximation, a non-covariant three-vector representation where all particles are confined to their mass shells and energy need not be conserved at individual vertices. An intuitive understanding of the processes embedded in the effective current density can be obtained by noting that the operator A is always associated with transitions from the $\eta-$space into the $\lambda-$space, so that with our present definitions, A serves to create meson+resonance+anti-nucleon existing states and $A^\dagger$ serves to restore purely two-nucleon states. In the second-order expansion of this current we observe direct terms J, initial- and final-state interaction terms $JA + A^\dagger J$, meson-recoil terms $A^\dagger JA$, and wave function re-orthonormalization terms $JA^\dagger A$ and $A^\dagger AJ$. The wave function re-orthonormalization terms result directly from the requirement that the transformation used to obtain eq. (2.14) is unitary - or equivalently, from the fact that we insist upon working with orthonormal wave functions. This point has been discussed in considerable detail elsewhere [3].

We select a momentum-space representation and perform a t-matrix expansion of the two-nucleon wave functions according to the standard procedure. With the photon field quantized in the Gupta-Bleuler formalism, we require only the transverse polarization vectors. Since these have a vanishing time component, we require only the spacial parts of the effective current density, so that,

$$\mathcal{M}_{fi} = N\vec{\epsilon}(\vec{k},\lambda)\langle \vec{p}_3\widetilde{\vec{p}_4;\alpha_f}|\vec{J}_{\text{eff}}(0)|\vec{p}_1\widetilde{\vec{p}_2;\alpha_i}\rangle \tag{2.22a}$$

$$+ N\vec{\epsilon}(\vec{k},\lambda)\langle \vec{p}_3\widetilde{\vec{p}_4;\alpha_f}|\vec{J}_{\text{eff}}(0)G_i t^{(+)}|\vec{p}_1\widetilde{\vec{p}_2;\alpha_i}\rangle \tag{2.22b}$$

$$+ N\vec{\epsilon}(\vec{k},\lambda)\langle \vec{p}_3\widetilde{\vec{p}_4;\alpha_f}|t^{(-)\dagger}G_f\vec{J}_{\text{eff}}(0)|\vec{p}_1\widetilde{\vec{p}_2;\alpha_i}\rangle \tag{2.22c}$$

$$+ N\vec{\epsilon}(\vec{k},\lambda)\langle \vec{p}_3\widetilde{\vec{p}_4;\alpha_f}|t^{(-)\dagger}G_f\vec{J}_{\text{eff}}(0)G_i t^{(+)}|\vec{p}_1\widetilde{\vec{p}_2;\alpha_i}\rangle\Big\} \tag{2.22d}$$

where G_i and G_f are η-space Green's functions describing the propagation of two-nucleon states and

$$N = -(2\pi)^6 m^{-2}\sqrt{E_1E_2E_3E_4} \tag{2.23}$$

The four terms shown in eq. (2.22) will be referred to as 'direct', 'initial-state', 'final-state' and 'rescattering' amplitudes respectively.

III. MODEL DEFINITION AND CALCULATION DETAILS

A. The RuhrPot Lagrangian

We adopt the strong-interaction Lagrangian densities,

$$\begin{aligned}
\mathcal{L}_{\text{NN}\pi} &= -ig_{\text{NN}\pi}\bar{\psi}[\lambda\gamma^5 - (1-\lambda)\frac{1}{2m}\gamma^5\gamma^\mu(i\partial_\mu)]\psi\vec{\pi}.\vec{\tau}\\
\mathcal{L}_{\text{NN}\eta} &= -\frac{g_{\text{NN}\eta}}{2m}\bar{\psi}\gamma^5\gamma^\mu\partial_\mu\psi\eta\\
\mathcal{L}_{\text{NN}\rho} &= -g_{\text{NN}\rho}\bar{\psi}\gamma^\mu\psi\vec{\rho}_\mu.\vec{\tau}\\
\mathcal{L}_{\text{NN}\omega} &= -g_{\text{NN}\omega}\bar{\psi}\gamma^\mu\psi\omega_\mu\\
\mathcal{L}_{\text{NN}\delta} &= -g_{\text{NN}\delta}\bar{\psi}\psi\vec{\delta}.\vec{\tau}\\
\mathcal{L}_{\text{NN}\epsilon} &= -g_{\text{NN}\epsilon}\bar{\psi}\psi\epsilon\\
\mathcal{L}_{\text{N}\Delta\pi} &= -\frac{g_{\text{N}\Delta\pi}}{2m}\bar{\psi}^\mu\vec{\tau}_{\text{N}\Delta}\psi\partial_\mu\vec{\pi} + \text{h.c.}\\
\mathcal{L}_{\text{N}\Delta\rho} &= -i\frac{g_{\text{N}\Delta\rho}}{2m}\bar{\psi}^\mu\gamma^5\gamma^\nu\vec{\tau}_{\text{N}\Delta}\psi\vec{\rho}_{\mu\nu} + \text{h.c.}
\end{aligned} \tag{3.1}$$

where $\rho_{\mu\nu} = \partial_\mu\rho_\nu - \partial_\nu\rho_\mu$, and we have denoted $g_{\text{N}\Delta\rho}{=}\mu_{\text{N}\Delta}g_\rho$, $g_{\text{NN}\rho}{=}\frac{1}{2}g_\rho$ and $g_{\text{NN}\omega}{=}\frac{1}{2}g_\omega$, where g_ρ and g_ω the strong charges for the ρ and ω gauge fields. For the electromagnetic-interaction we use the Lorentz-Heaviside system with natural units, where the charge of the proton is defined as $e_p = +\sqrt{4\pi\alpha}$ with $\alpha \sim 1/137.04$, and we adopt the Lagrangians,

$$\begin{aligned}
\mathcal{L}_{\text{NN}\gamma} &= -e_{\text{N}}\bar{\psi}\gamma^\mu\psi\mathcal{A}_\mu + \frac{e_p\kappa_{\text{N}}}{2m}\bar{\psi}\sigma^{\mu\nu}\partial_\nu\mathcal{A}_\mu\psi\\
\mathcal{L}_{\text{PV}\gamma} &= -\frac{e_p g_{\text{PV}\gamma}}{2m_{\text{V}}}\epsilon^{\mu\nu\sigma\tau}F_{\mu\nu}\vec{\phi}^{\text{V}}_\sigma.\partial_\tau\vec{\phi}^{\text{P}}\\
\mathcal{L}_{\text{N}\Delta\gamma} &= -i\frac{e_p}{2m}\mu_{\text{N}\Delta}\bar{\psi}^\mu\gamma^5\gamma^\nu\tau^3_{\text{N}\Delta}\psi F_{\mu\nu} + \text{h.c.}
\end{aligned} \tag{3.2}$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, $\vec{\phi}^{\text{V}}{=}\vec{\rho}$ or ω, $\vec{\phi}^{\text{P}}{=}\vec{\pi}$ or η, $e_{\text{N}}{=}e_p\frac{(1+\tau^3)}{2}$ and $\mu_{\text{N}}{=}\frac{(1+\kappa^{\text{is}})}{2} + \frac{(1+\kappa^{\text{iv}})}{2}\tau^3 = 1{+}\kappa_{\text{N}}$ with $\kappa^{\text{is}}{=}{-0.12}$ and $\kappa^{\text{iv}}{=}3.706$.

These Lagrangians describe the bare coupling of mass renormalized fields. The form factors describing the coupling constant renormalization have been calculated [23] as a coupled set of integral equations yielding results which can be accurately parameterized as monopoles with the regularization scales shown in Table 1. These renormalized couplings are implemented by replacing the Lagrangians with the vertex functions describing their dressed counterparts,

$$\begin{aligned}
\Gamma_{\text{NN}\pi} &= -ig_{\text{NN}\pi}F_{\text{NN}\pi}\bar{\psi}[\lambda\gamma^5 - (1-\lambda)\frac{1}{2m}\gamma^5\gamma^\mu(i\partial_\mu)]\psi\vec{\pi}.\vec{\tau}\\
\Gamma_{\text{NN}\eta} &= -\frac{g_{\text{NN}\eta}F_{\text{NN}\eta}}{2m}\bar{\psi}\gamma^5\gamma^\mu\partial_\mu\psi\eta\\
\Gamma_{\text{NN}\rho} &= -g_{\text{NN}\rho}\bar{\psi}[F^{(1)}_{\text{NN}\rho}\gamma^\mu - \frac{\kappa_\rho F^{(2)}_{\text{NN}\rho}}{2m}\sigma^{\mu\nu}\partial_\nu]\psi\vec{\rho}_\mu.\vec{\tau}\\
\Gamma_{\text{NN}\omega} &= -g_{\text{NN}\omega}\bar{\psi}[F^{(1)}_{\text{NN}\omega}\gamma^\mu - \frac{\kappa_\omega F^{(2)}_{\text{NN}\omega}}{2m}\sigma^{\mu\nu}\partial_\nu]\psi\omega_\mu\\
\Gamma_{\text{NN}\delta} &= -g_{\text{NN}\delta}F_{\text{NN}\delta}\bar{\psi}\psi\vec{\delta}.\vec{\tau}\\
\Gamma_{\text{NN}\epsilon} &= -g_{\text{NN}\epsilon}F_{\text{NN}\epsilon}\bar{\psi}\psi\epsilon\\
\Gamma_{\text{N}\Delta\pi} &= -\frac{g_{\text{N}\Delta\pi}F_{\text{N}\Delta\pi}}{2m}\bar{\psi}^\mu\vec{\tau}_{\text{N}\Delta}\psi\partial_\mu\vec{\pi} + \text{h.c.}
\end{aligned}$$

$$\Gamma_{\mathrm{N}\Delta\rho} = -i\frac{g_{\mathrm{NN}\rho}G_{\mathrm{N}\Delta\rho}}{2m}\frac{g_{\mathrm{N}\Delta\pi}}{g_{\mathrm{NN}\pi}}\bar{\psi}^{\mu}\gamma^{5}\gamma^{\nu}\vec{\tau}_{\mathrm{N}\Delta}\psi\vec{\rho}_{\mu\nu} + \mathrm{h.c.} \quad (3.3)$$

where $G_{\mathrm{N}\Delta\rho}=F^{(1)}_{\mathrm{N}\Delta\rho}+\kappa_\rho F^{(2)}_{\mathrm{N}\Delta\rho}$ and we normalize all form factors as $F(0) = 1$. As in other exchange current applications [37] the experimentally unknown values of $g_{\mathrm{N}\Delta\rho} = 20.73$ and $\mu_{\mathrm{N}\Delta}$=3.993 are fixed according to SU(6) [38,39] and vector-meson dominance as,

$$\mu_{\mathrm{N}\Delta} = \mu^{\mathrm{iv}}_{\mathrm{N}}\frac{g_{\mathrm{N}\Delta\pi}}{g_{\mathrm{NN}\pi}} \qquad g_{\mathrm{N}\Delta\rho} = g_{\mathrm{NN}\rho}(1+\kappa_\rho)\frac{g_{\mathrm{N}\Delta\pi}}{g_{\mathrm{NN}\pi}} \quad (3.4)$$

where $\mu^{\mathrm{iv}}_{\mathrm{N}} = \frac{1}{2}G^{\mathrm{V}}_{\mathrm{M}}(0) = 2.353$. Note that the tensor couplings κ_ρ and κ_ω are absent from the Lagrangians of eq. (3.1) but appear in the dressed vertex functions of eq. (3.3) since they are directly computed from the loop-integrals appearing in the form factor calculation [23]. A similar consideration for the electromagnetic form factors is obviously not required for the real photon.

The NNα properties are taken from the fit of the RuhrPot two-nucleon interaction [28] to the NN-scattering data. The form factor scales adopted in ref [28] were actually calculated within a non-relativistic framework [40,41], but the recent relativistic calculation [23] has confirmed the parameterizations. We acknowledge some ambiguity in the signs of the PVγ coupling constants [42,43] but adopt $g_{\rho\pi\gamma}$=0.53, $g_{\omega\pi\gamma}$=2.58, $g_{\rho\eta\gamma}$=1.39 and $g_{\omega\eta\gamma}$=0.15, as reported in ref [44]. We use the experimental result $g_{\mathrm{N}\Delta\pi} = 28.85$, which is consistent with the Chew-Low [45] and strong-coupling [46] models and, moreover, with the form factor calculations [23]. We will not fiddle with these values in order to optimize selected experimental results since this would spoil the consistency between the calculation of the meson-baryon form-factors, the NN-interaction and the exchange currents.

β	m_β (MeV)	$g_{\mathrm{NN}\beta}$	κ_β (GeV)$^{-2}$	Σ_β	κ_{Σ_β}	$g_{\Delta\mathrm{N}\beta}$
π	136.5	12.922	-	49.516	-	28.85
η	548.8	6.015	-	-	-	-
ρ	776.9	1.651	6.400	0.0124	28.105	20.73
ω	782.4	4.945	1.088	12.379	0.4334	-
δ	983.0	6.043	-	-	-	-
ϵ	975.0	10.567	-	5.6911	-	-

TABLE I. RuhrPot parameters adopted in the present calculation. All NN-meson form factors are taken from direct calculation. For the ϵ meson this requires a meson scale of Λ_1=0.6 GeV, whereas all other mesons require Λ_1=0.8 GeV. (Further details can be found in [23,28]). We adopt the experimental results $g_{\rho\pi\gamma}$=0.53, $g_{\omega\pi\gamma}$=2.58, $g_{\rho\eta\gamma}$=1.39, $g_{\omega\eta\gamma}$=0.15 and κ^{is}=-0.12, κ^{iv}=3.706. SU(6) and vector dominance indicate $\mu_{\mathrm{N}\Delta}$=3.993.

B. Impulse and Exchange Currents

We describe here the impulse and meson-exchange currents J_{eff}, as required in eq. (2.22). We adopt a partition of Hilbert spaces into meson+resonance+anti-nucleon - vacuum and -existing parts, as described in section II. In the present work we confine our attention to leading-order exchange currents involving the electromagnetic coupling to the NN, $\bar{\mathrm{N}}$N, PV=$\rho\pi$, $\omega\pi$, $\rho\eta$ and $\omega\eta$ and NΔ currents, so that,

$$J_{\mathrm{eff}} = J^{\mathrm{NN}}_{\mathrm{eff}} + J^{\bar{\mathrm{N}}\mathrm{N}}_{\mathrm{eff}} + J^{\mathrm{PV}}_{\mathrm{eff}} + J^{\mathrm{N}\Delta}_{\mathrm{eff}} \quad (3.5)$$

The effective current can therefore be derived unambiguously from eqs. (3.2), (3.3), (2.12), (2.18-2.19) and (2.21). Throughout we describe the momenta of a meson with mass m_β with,

$$\vec{q}_1 = \vec{p}_3 - \vec{p}_1, \;\; \omega_1 = \sqrt{\vec{q}_1^{\,2} + m_\beta^2}, \;\; q_1^0 = E_3 - E_1, \;\; Q_1^2 = -q_1^2$$
$$\vec{q}_2 = \vec{p}_4 - \vec{p}_2, \;\; \omega_2 = \sqrt{\vec{q}_2^{\,2} + m_\beta^2}, \;\; q_2^0 = E_4 - E_2, \;\; Q_2^2 = -q_2^2 \quad (3.6)$$

and, denoting the nucleon and Δ-isobar masses as m and m_Δ, we condense our notation with,

$$\vec{p}_{ik} = \begin{cases} \vec{p}_i - \vec{k} & \text{for } i\text{=1, 2} \\ \vec{p}_i + \vec{k} & \text{for } i\text{=3, 4} \end{cases}$$
$$E_{ik} = \sqrt{\vec{p}_{ik} + m^2}, \qquad \mathcal{E}_{ik} = E_{ik} + m,$$
$$E_{\Delta ik} = \sqrt{\vec{p}_{ik} + m_\Delta^2}, \qquad \mathcal{E}_{\Delta ik} = E_{\Delta ik} + m_\Delta \quad (3.7)$$

1. *Impulse and Exchange currents with the Relativistic NNγ vertex*

For the partition of Hilbert spaces defined in section II B, all contributions involving a vertex where the photon couples to the nucleon current must satisfy $\lambda J_{\mathrm{NN}}\eta = \eta J_{\mathrm{NN}}\lambda = 0$, so that eq. (3.5) requires,



FIG. 4. NNγ wave function re-orthonormalization and meson-recoil exchange currents. These currents are necessary to preserve the orthonormality of the initial- and final-state wave functions.

$$
\begin{aligned}
\langle \vec{p}_3\vec{p}_4|\left[J^{\mathrm{NN}}_{\mathrm{eff}}\right]^{\mu}|\vec{p}_1\vec{p}_2\rangle = {} & J^{\mu}_{\mathrm{NN}\gamma}[1](\vec{p}_1,\vec{p}_3)\delta(\vec{p}_4-\vec{p}_2)\\
& + J^{\mu}_{\mathrm{NN}\gamma}[2](\vec{p}_2,\vec{p}_4)\delta(\vec{p}_3-\vec{p}_1)\\
& + \langle \vec{p}_3\vec{p}_4|J^{\mu}_{\mathrm{wfrr}}|\vec{p}_1\vec{p}_2\rangle
\end{aligned}
\tag{3.8}
$$

where the first two terms describe the impulse currents for nucleons 1 and 2, and the last term denotes the wave function re-orthonormalization and meson recoil currents,

$$
\begin{aligned}
&\langle \vec{p}_3\vec{p}_4|J^{\mu}_{\mathrm{wfrr}}|\vec{p}_1\vec{p}_2\rangle = -g_{\sigma\tau}\sum_{\beta}\Big[\\
&D^{\mathrm{NN}\beta}_{\mathrm{abc}}J^{\mu}_{\mathrm{NN}\gamma}[1](\vec{p}_{3k},\vec{p}_3)H^{\sigma}_{\mathrm{NN}\beta}[1](\vec{p}_1,\vec{p}_{3k})H^{\tau}_{\mathrm{NN}\beta}[2](\vec{p}_2,\vec{p}_4)\\
&+D^{\mathrm{NN}\beta}_{\mathrm{def}}H^{\sigma}_{\mathrm{NN}\beta}[1](\vec{p}_{1k},\vec{p}_3)H^{\tau}_{\mathrm{NN}\beta}[2](\vec{p}_2,\vec{p}_4)J^{\mu}_{\mathrm{NN}\gamma}[1](\vec{p}_{1k},\vec{p}_1)\Big]\\
&+(1,3\rightleftharpoons 2,4)
\end{aligned}
\tag{3.9}
$$

where $\beta=\pi$, η, ρ, ω, δ and ϵ, and the factor $-g_{\sigma\tau}$ resulting from the contraction of the vector meson polarizations and all references to the Lorentz indices σ and τ are to be ignored for the scalar mesons. Explicit expressions for the vertex functions $H_{\mathrm{NN}\beta}$ and current $J_{\mathrm{NN}\gamma}$ are supplied in app. B. The propagator functions are labeled in correspondence to Fig. 4 and are defined as,

$$
\begin{aligned}
D^{\beta}_{\mathrm{abc}} = {} & \frac{-\frac{1}{2}}{[E_3+E_4-E_{3k}-E_2-\omega_2][E_1-E_{3k}-\omega_2]}\\
& + \frac{1}{[E_4-E_2-\omega_2][E_1-E_{3k}-\omega_2]}\\
& + \frac{-\frac{1}{2}}{[E_3-E_1-\omega_2][E_2-E_4-\omega_2]}\\
D^{\beta}_{\mathrm{def}} = {} & \frac{-\frac{1}{2}}{[E_3-E_{1k}-\omega_2][E_1+E_2-E_{1k}-E_4-\omega_2]}\\
& + \frac{1}{[E_3-E_{1k}-\omega_2][E_2-E_4-\omega_2]}\\
& + \frac{-\frac{1}{2}}{[E_4-E_2-\omega_2][E_1-E_3-\omega_2]}
\end{aligned}
\tag{3.10}
$$

In the static limit we note that $D^{\beta}_{\mathrm{abc}} = D^{\beta}_{\mathrm{def}} = 0$ so that the wave function renormalization and meson recoil exchange currents J^{μ}_{wfrr} simply vanish. The same conclusion can be reached within the soft-photon approximation in the barycentric frame.

Since we will avoid these approximations, we are forced to accept that a relativistic description of the photon coupling to the positive-frequency components of the impulse current $J^{\mu}_{\mathrm{NN}\gamma}$ necessarily leads to an effective current density J_{eff} comprising both one- and two-body operators. These two-body contributions have, to date, never been explicitly included in any of the bremsstrahlung calculations that seek to include the relativistic components of the NNγ vertex.

2. Pair Currents

The sum of one-body impulse-currents $J^{\mu}_{\mathrm{NN}\gamma}$ and the wave function re-orthonormalization and meson-recoil exchange currents J^{μ}_{wfrr} do not exhaust the requirements needed to obtain a relativistic description of the photon coupling to the nucleon current density since the off-shell nucleon comprises a linear super-position of positive and (so far neglected) negative frequency components. In the Feynman-Stückelberg approach, the negative-frequency components of the off-shell nucleon field are interpreted as anti-particles, so we are led to introduce the photon coupling to the N$\bar{\mathrm{N}}$-pair creation and annihilation currents.

Within our partition of Hilbert spaces, the photon coupling to the N$\bar{\mathrm{N}}$-pair creation and annihilation vertices must satisfy $\eta J_{\bar{N}N}\eta = 0$, so that eq. (3.5) requires,

$$
\begin{aligned}
&\langle \vec{p}_3\vec{p}_4|\left[J^{\mathrm{N\bar{N}}}_{\mathrm{eff}}\right]^{\mu}|\vec{p}_1\vec{p}_2\rangle = -g_{\sigma\tau}\sum_{\beta}\Big[\\
&D^{\mathrm{N\bar{N}}\beta}_{\mathrm{abc}}H^{\sigma}_{\mathrm{N\bar{N}}\beta}[1](\vec{p}_1,\vec{p}_{3k})H^{\tau}_{\mathrm{NN}\beta}[2](\vec{p}_2,\vec{p}_4)J^{\mu}_{\mathrm{N\bar{N}}\gamma}[1](\vec{p}_{3k},\vec{p}_3)\\
&+D^{\mathrm{N\bar{N}}\beta}_{\mathrm{def}}J^{\mu}_{\mathrm{N\bar{N}}\gamma}[1](\vec{p}_{1k},\vec{p}_1)H^{\sigma}_{\mathrm{N\bar{N}}\beta}[1](\vec{p}_{1k},\vec{p}_3)H^{\tau}_{\mathrm{NN}\beta}[2](\vec{p}_2,\vec{p}_4)\Big]\\
&+(1,3\rightleftharpoons 2,4)
\end{aligned}
\tag{3.11}
$$



FIG. 5. $\mathrm{N\bar{N}}$ pair creation and annihilation meson-exchange currents. These currents are necessary for a relativistic description of the $\mathrm{NN}\gamma$ vertex.

where $\beta = \pi$, η, ρ, ω, δ and ϵ, and the factor $-g_{\sigma\tau}$ and all references to the Lorentz indices σ and τ are to be ignored for the scalar mesons. Explicit expressions for the pair-creation and -annihilation currents $J_{\mathrm{N\bar{N}}\gamma}$ and $J_{\mathrm{\bar{N}N}\gamma}$ and all required vertex functions are described in app. B.

The propagator functions are labeled in correspondence to Fig. 5 and are defined as,

$$
\begin{aligned}
D^{\beta}_{\mathrm{abc}} = & \frac{1}{[-E_{3k} - E_1 - \omega_2][E_4 - E_{3k} - E_1 - E_2]} \\
& + \frac{1}{[E_4 - E_2 - \omega_2][E_4 - E_{3k} - E_1 - E_2]} \\
& + \frac{1}{[-E_{3k} - E_1 - \omega_2][E_2 - \omega_2 - E_4]} \\
D^{\beta}_{\mathrm{def}} = & \frac{1}{[E_2 - E_{1k} - E_3 - E_4][-E_{1k} - E_3 - \omega_2]} \\
& + \frac{1}{[E_2 - E_{1k} - E_3 - E_4][E_2 - \omega_2 - E_4]} \\
& + \frac{1}{[E_4 - \omega_2 - E_2][-E_{1k} - E_3 - \omega_2]}
\end{aligned}
\tag{3.12}
$$

We will adopt these expressions for our numerical work. Nonetheless, it is interesting to consider the corresponding result under various approximations. If we demand energy conservation across the current matrix elements, then eq. (3.12) reduces to,

$$
\begin{aligned}
D^{\beta}_{\mathrm{abc}} &= \frac{-2\omega_2}{(E_3+k+E_{3k})(q_2^2-m_\beta^2)} \overset{\mathrm{nr}}{\sim} \frac{2}{\omega_2(2m+k)} \overset{\mathrm{spa}}{\sim} \frac{1}{\omega_2 m} \\
D^{\beta}_{\mathrm{def}} &= \frac{-2\omega_2}{(E_1-k+E_{1k})(q_2^2-m_\beta^2)} \overset{\mathrm{nr}}{\sim} \frac{2}{\omega_2(2m-k)} \overset{\mathrm{spa}}{\sim} \frac{1}{\omega_2 m}
\end{aligned}
\tag{3.13}
$$

where we also provide the static-limit and soft-photon reductions. The corresponding π-exchange contribution under such approximations are,

$$
\begin{aligned}
\langle \vec{p}_3\vec{p}_4 | \left[J^{\mathrm{\bar{N}N}}_{\mathrm{eff}}\right]^{\mu} | \vec{p}_1\vec{p}_2 \rangle &\overset{\mathrm{nr}}{\sim} \frac{\lambda e_p g^2_{\mathrm{NN}\pi} F^2_{\mathrm{NN}\pi}(\vec{q}_2)}{(2\pi)^6 4m(\vec{q}_2^{\,2} + m_\pi^2)} \vec{\sigma}_1(\vec{\sigma}_2.\vec{q}_2) \left\{ \frac{(1+\tau_1^z)\vec{\tau}_1.\vec{\tau}_2}{(2m-k)} - \frac{\vec{\tau}_1.\vec{\tau}_2(1+\tau_1^z)}{(2m+k)} \right\} + (1 \rightleftharpoons 2) \\
\langle \vec{p}_3\vec{p}_4 | \left[J^{\mathrm{\bar{N}N}}_{\mathrm{eff}}\right]^{\mu} | \vec{p}_1\vec{p}_2 \rangle &\overset{\mathrm{spa}}{\sim} \frac{-\lambda e_p g^2_{\mathrm{NN}\pi} F^2_{\mathrm{NN}\pi}(\vec{q}_2)}{(2\pi)^6 4m^2(\vec{q}_2^{\,2} + m_\pi^2)} \vec{\sigma}_1(\vec{\sigma}_2.\vec{q}_2)(i\vec{\tau}_1 \times \vec{\tau}_2)^3 + (1 \rightleftharpoons 2)
\end{aligned}
\tag{3.14}
$$

Both of these results scale linearly with the parameter λ controlling the admixture of ps- and pv-couplings in the $\mathrm{NN}\pi$ Lagrangian, but the well-known isovector structure of the non-relativistic pair currents holds only for soft photons. Since the data with which we will compare our results was planned to maximize the photon energy, we anticipate a non-negligible contribution fom the isoscalar components of eq (3.11). This offers the possibility of studying λ without the complications of describing the many leading-order exchange currents that contribute to np-bremsstrahlung.

3. PVγ Currents

The ω meson (782.4 MeV) decays as $\omega \rightarrow \pi^0\gamma$ with an 8.7% branching ratio and indicates the coupling constant of $g_{\omega\pi\gamma}$=2.58. As such, the $\omega\pi\gamma$ exchange currents can be expected to make a non-trivial contribution to both



FIG. 6. $\mathrm{VP}\gamma = \rho\pi\gamma + \omega\pi\gamma + \rho\eta\gamma + \omega\eta\gamma$ exchange currents. When energy is conserved across these current matrix elements, the time-ordered graphs exactly sum to form the corresponding Feynman diagrams.

the pp- and np-bremsstrahlung observables. Analogous arguments indicate that the $\rho\pi\gamma$ contributions will be large in np-bremsstrahlung, and perhaps also of some lesser importance in pp-bremsstrahlung.

Our desire to preserve complete consistency between the form factors, NN-interaction and the exchange currents leads us to introduce all leading-order exchange currents describing the photon coupling to the decay of all vector-meson mesons present in the form factor and NN-interaction calculations. We therefore include the $\mathrm{PV}\gamma = \rho\pi\gamma + \omega\pi\gamma + \rho\eta\gamma + \omega\eta\gamma$ exchange currents as shown in Fig. 6. Each of these vertices satisfies $\eta J_{\mathrm{P\bar{V}}}\eta = 0$, so that after making use of eqs. (3.2), (2.12) and (2.21), eq. (3.5) requires,

$$\begin{aligned}\langle\vec{p}_3\vec{p}_4|\left[J^{\mathrm{VP}}_{\mathrm{eff}}\right]^{\mu}|\vec{p}_1\vec{p}_2\rangle &= \frac{\sqrt{4\omega_{\mathrm{P}}(\vec{q}_1)\omega_{\mathrm{V}}(\vec{q}_2)}}{(2\pi)^3 m_{\mathrm{V}}[q_1^2 - m_{\mathrm{P}}^2][q_2^2 - m_{\mathrm{V}}^2]}\\ &\times H_{\mathrm{NNP}}[1](\vec{p}_1,\vec{p}_3)\epsilon_{\nu}(\hat{q}_2,\lambda_{\mathrm{V}})H^{\nu}_{\mathrm{NNV}}[2](\vec{p}_2,\vec{p}_4)J^{\mu}_{\mathrm{VP}\gamma}(q_1,q_2)\\ &+(1,3\rightleftharpoons 2,4) \qquad (3.15)\end{aligned}$$

where $J_{\mathrm{VP}\gamma}$ is given in App. B. We require only the 3-vector current, for which the relativistic form is,

$$\begin{aligned}\langle\vec{p}_3\vec{p}_4|\vec{J}^{\mathrm{VP}}_{\mathrm{eff}}|\vec{p}_1\vec{p}_2\rangle &= -\frac{e_p g_{\mathrm{VP}\gamma}F_{\mathrm{VP}\gamma}\sqrt{4\omega_{\mathrm{P}}(\vec{q}_1)\omega_{\mathrm{V}}(\vec{q}_2)}}{(2\pi)^3 m_{\mathrm{V}}[q_1^2 - m_{\mathrm{P}}^2][q_2^2 - m_{\mathrm{V}}^2]}\\ &\times H_{\mathrm{NNP}}[1](\vec{p}_1,\vec{p}_3)\Big\{H^0_{\mathrm{NNV}}[2](\vec{p}_2,\vec{p}_4)[\vec{q}_1\times\vec{q}_2]\\ &\quad+\vec{H}_{\mathrm{NNV}}[2](\vec{p}_2,\vec{p}_4)\times[(q_2)_0\vec{q}_1 - (q_1)_0\vec{q}_2]\Big\}\\ &+(1,3\rightleftharpoons 2,4) \qquad (3.16)\end{aligned}$$

We will not resort to the non-relativistic limit, but we realize it implies $q_1^0{=}q_2^0{=}0$, so that we recover the well-known result for the emission of a real photon of momentum $\vec{k}$,

$$\begin{aligned}\langle\vec{p}_3\vec{p}_4|\vec{J}^{\mathrm{VP}}_{\mathrm{eff}}|\vec{p}_1\vec{p}_2\rangle &\overset{\mathrm{nr}}{\sim} \frac{ie_p g_{\mathrm{VP}\gamma}g_{\mathrm{NNV}}g_{\mathrm{NNP}}}{(2\pi)^6 2m_{\mathrm{V}}m[\vec{q}_1^{\,2}+m_{\mathrm{P}}^2][\vec{q}_2^{\,2}+m_{\mathrm{V}}^2]}\\ &\times(\vec{\sigma}_1.\vec{q}_1)(\vec{q}_1\times\vec{q}_2)(\vec{\tau}_1)_{\mathrm{P}}(\vec{\tau}_2)_{\mathrm{V}}+(1,3\rightleftharpoons 2,4) \qquad (3.17)\end{aligned}$$

where $(\vec{\tau}_1)_{\mathrm{P}}(\vec{\tau}_2)_{\mathrm{V}} = \vec{\tau}_1.\vec{\tau}_2$, $\vec{\tau}_1^{\,0}$, $\vec{\tau}_2^{\,0}$ and $\mathbb{1}$ for the $\rho\pi\gamma$, $\omega\pi\gamma$, $\rho\eta\gamma$ and $\omega\eta\gamma$ currents respectively.

4. NΔγ Exchange Currents

All contributions involving a vertex where the photon couples to the NΔ current must satisfy $\eta J_{\Delta\mathrm{N}}\eta = \eta J_{\mathrm{N}\Delta}\eta = 0$, so that eq. (3.5) requires,

$$\begin{aligned}\langle\vec{p}_3\vec{p}_4|\vec{J}^{\mathrm{N}\Delta}_{\mathrm{eff}}|\vec{p}_1\vec{p}_2\rangle &= -g_{\sigma\tau}\sum_{\beta}\Big[\\ &D^{\beta}_{\mathrm{abc}}\vec{J}_{\mathrm{N}\Delta\gamma}[1](\vec{p}_{3k},\vec{p}_3)H^{\sigma}_{\Delta\mathrm{N}\beta}[1](\vec{p}_1,\vec{p}_{3k})H^{\tau}_{\mathrm{NN}\beta}[2](\vec{p}_2,\vec{p}_4)\\ &+D^{\beta}_{\mathrm{def}}H^{\sigma}_{\mathrm{N}\Delta\beta}[1](\vec{p}_{1k},\vec{p}_3)\vec{J}_{\Delta\mathrm{N}\gamma}[1](\vec{p}_1,\vec{p}_{1k})H^{\tau}_{\mathrm{NN}\beta}[2](\vec{p}_2,\vec{p}_4)\Big]\\ &+(1,3\rightleftharpoons 2,4) \qquad (3.18)\end{aligned}$$

where $\beta{=}\vec{\pi}$ or $\vec{\rho}$ and the factor $-g_{\sigma\tau}$ resulting from the contraction of the ρ-meson polarizations and all references to the Lorentz indices σ and τ are to be ignored for $\beta{=}\pi$. Explicit expressions for the vertex functions and currents shown in eq. (3.18) are supplied in app. B. We ignore all negative frequency resonance contributions. The propagator functions are labeled in correspondence to Fig. 7 and are defined in analogy to the previous sections. Using energy conservation for the current matrix elements and introducing the Δ decay width Γ_{Δ} via [47,48] $E_{\Delta k}\to E_{\Delta k} - i\Gamma_{\Delta}/2$, the propagators reduce to,

$$\begin{aligned}D^{\beta}_{\mathrm{abc}} &= \frac{2\omega_2}{(q_2^2 - m_{\beta}^2)(E_3 + k - E_{\Delta 3k} + i\Gamma_{\Delta}/2)}\\ D^{\beta}_{\mathrm{def}} &= \frac{2\omega_2}{(q_2^2 - m_{\beta}^2)(E_1 - k - E_{\Delta 1k} + i\Gamma_{\Delta}/2)} \qquad (3.19)\end{aligned}$$

An exact calculation of eq. (3.18) can be achieved with the use of the vertex functions and currents given in App. B. However, at present there exists some uncertainty in the coupling constants $g_{\mathrm{N}\Delta\rho}$ and $\mu_{\mathrm{N}\Delta}$, so that such a rigorous procedure is of limited interest. We simplify matters by dropping terms of order $p^2/(E+m)^2$



FIG. 7. N$\Delta\gamma$ $\pi-$ and $\rho-$meson exchange currents. In the RuhrPot model the $\pi-$exchange contributions represent the largest of the two-body currents in pp-bremsstrahlung but the $\rho-$exchange contributions are very small. When energy is conserved across these current matrix elements, the time-ordered graphs exactly sum to form the corresponding Feynman diagrams.

beyond leading order - an approximation which does not involve any $p/m-$expansion and should be accurate to within a few per-cent at energies below the π-production threshold. This is surely adequate for the first-order S_{fi}-matrix described in sec II A.

We introduce the condensed notation,

$$\vec{\mathcal{P}}_i = \frac{\vec{p}_i}{\mathcal{E}_i}, \quad \vec{\mathcal{P}}_{\Delta ki} = \frac{\vec{p}_{ik}}{\mathcal{E}_{\Delta ik}}, \quad \vec{\mathcal{Q}}_{ij} = \vec{p}_{ik}\left(\frac{E_j}{m_\Delta}\right) - \vec{p}_j,$$
$$\vec{\mathcal{K}}_i = \vec{p}_{ik}\Big(\frac{2E_{\Delta ik} - E_i}{m_\Delta}\Big) - \vec{p}_i,$$
$$\vec{\mathcal{L}}_i = \left(\frac{\vec{p}_{ik}}{\mathcal{E}_{\Delta ik}} - \frac{\vec{p}_i}{\mathcal{E}_i}\right)(m + m_\Delta) \qquad i = 1,2,3,4. \tag{3.20}$$

and proceed to calculate separately the contributions from π- and ρ-exchange,

$$\vec{J}^{\mathrm{N}\Delta}_{\mathrm{eff}} = \vec{J}^{\mathrm{N}\Delta}_{\mathrm{eff}}(\pi) + \vec{J}^{\mathrm{N}\Delta}_{\mathrm{eff}}(\rho) \tag{3.21}$$

5. N$\Delta\gamma$ $\pi-$Exchange Currents

The effective current describing the excitation of intermediate Δ-isobars through π-exchange can now be obtained from eqs. (3.18) and App. B. For the real photon, we require only the spacial current, which takes the form,

$$\begin{aligned}
&\langle\vec{p}_3\vec{p}_4|\vec{J}^{\mathrm{N}\Delta}_{\mathrm{eff}}(\pi)|\vec{p}_1\vec{p}_2\rangle \\
&= N_1\vec{\sigma}_2.(\vec{\mathcal{P}}_4-\vec{\mathcal{P}}_2)\Big[2\tau_2^3-(i\tau_1\times\tau_2)^3\Big] \\
&\quad\times\Big[(\vec{\mathcal{K}}_1+\vec{\mathcal{L}}_1)\times\vec{\mathcal{Q}}_{13}+(i\vec{\sigma}_1\times\vec{\mathcal{Q}}_{13})\times\vec{\mathcal{L}}_1 \\
&\quad+i\vec{\sigma}_1\times[(\vec{\mathcal{K}}_1-\vec{\mathcal{L}}_1)\times\vec{\mathcal{Q}}_{13}]-2i\vec{\sigma}_1[(\vec{\mathcal{K}}_1-\vec{\mathcal{L}}_1).\vec{\mathcal{Q}}_{13}]\Big] \\
&+ N_3\vec{\sigma}_2.(\vec{\mathcal{P}}_4-\vec{\mathcal{P}}_2)\Big[2\tau_2^3+(i\tau_1\times\tau_2)^3\Big] \\
&\quad\times\Big[(\vec{\mathcal{K}}_3+\vec{\mathcal{L}}_3)\times\vec{\mathcal{Q}}_{31}-(i\vec{\sigma}_1\times\vec{\mathcal{Q}}_{31})\times\vec{\mathcal{L}}_3 \\
&\quad-i\vec{\sigma}_1\times[(\vec{\mathcal{K}}_3-\vec{\mathcal{L}}_3)\times\vec{\mathcal{Q}}_{31}]+2i\vec{\sigma}_1[(\vec{\mathcal{K}}_3-\vec{\mathcal{L}}_3).\vec{\mathcal{Q}}_{31}]\Big] \\
&+ (1,3 \rightleftharpoons 2,4)
\end{aligned} \tag{3.22}$$

where,

$$N_0 = \frac{-ie_p\mu_{\mathrm{N}\Delta}g_{\mathrm{N}\Delta\pi}g_{\mathrm{NN}\pi}}{(2\pi)^6 72m^2}\left[\frac{\mathcal{E}_1\mathcal{E}_2\mathcal{E}_3\mathcal{E}_4}{16E_1E_2E_3E_4}\right]^{\frac{1}{2}}\frac{F_{\mathrm{NN}\pi}F_{\mathrm{N}\Delta\pi}}{\omega_\pi(\vec{q}_2^{\,2})},$$
$$N_1 = \frac{N_0\mathcal{E}_{\Delta 1k}D^\pi_{\mathrm{def}}}{2E_{\Delta 1k}}, \qquad N_3 = \frac{N_0\mathcal{E}_{\Delta 3k}D^\pi_{\mathrm{abc}}}{2E_{\Delta 3k}} \tag{3.23}$$

We will adopt eq. (3.22) for our numerical applications and use it to consider the merit of various approximations that are conventionally adopted to recover a simplified operator structure.

The first level of approximation involves taking the static limit and ignoring the N-Δ mass difference in eq. (3.20). We call this the *vertex static limit* approximation and note that it is equivalent to casting eq. (3.22) as,

$$\begin{aligned}
&\langle\vec{p}_3\vec{p}_4|\vec{J}^{\mathrm{N}\Delta}_{\mathrm{eff}}(\pi)|\vec{p}_1\vec{p}_2\rangle \\
&= (N_1+N_3)\frac{(\vec{\sigma}_2.\vec{q}_2)}{2m}\left[4\tau_2^3\vec{q}_2 - (i\vec{\tau}_1\times\vec{\tau}_2)^3(i\vec{\sigma}_1\times\vec{q}_2)\right]\times\vec{k} \\
&- 2(N_1-N_3)\frac{(\vec{\sigma}_2.\vec{q}_2)}{2m}\left[\tau_2^3(i\vec{\sigma}_1\times\vec{q}_2) - (i\vec{\tau}_1\times\vec{\tau}_2)^3\vec{q}_2\right]\times\vec{k} \\
&+(1,3 \rightleftharpoons 2,4)
\end{aligned} \tag{3.24}$$

The substantial simplification results primarily because the vertex static limit approximation indicates $\mathcal{K}_1{=}\mathcal{L}_1{=}-\vec{k}$, and $\mathcal{K}_3{=}\mathcal{L}_3{=}\vec{k}$, so that all operator structures involving $\mathcal{K}_i - \mathcal{L}_i$ immediately vanish. However, if we consider the static limit with the more reasonable approximation $m_\Delta \sim \frac{4}{3}m$, then we find

$$\mathcal{K}_1 + \mathcal{L}_1 \sim -\frac{17}{8}\vec{k} - \frac{1}{24}\vec{p}_1, \qquad \mathcal{K}_1 - \mathcal{L}_1 \sim -\frac{3}{8}\vec{k} + \frac{13}{24}\vec{p}_1$$
$$\mathcal{K}_3 + \mathcal{L}_3 \sim \frac{17}{8}\vec{k} - \frac{1}{24}\vec{p}_3, \qquad \mathcal{K}_3 - \mathcal{L}_3 \sim \frac{3}{8}\vec{k} + \frac{13}{24}\vec{p}_3 \tag{3.25}$$

so that the $\mathcal{K}_i + \mathcal{L}_i$ terms surviving in the vertex static limit are rather well approximated, but the neglected $\mathcal{K}_i - \mathcal{L}_i$ terms are poorly represented.

The second level of approximation involves casting the complete expression in the static limit. This is the approximation we adopted in earlier work [1,15]. At the small momentum transfers relevant to the present numerical application, this *complete static limit* approximation will differ little from the vertex static limit and, if we further set $\Gamma_\Delta = 0$ and drop the form factor dependence, it is equivalent to casting eq. (3.22) in the simpler form,

$$
\begin{aligned}
\langle \vec{p}_3\vec{p}_4 | \vec{J}^{\rm N\Delta}_{\rm eff}(\pi) | \vec{p}_1\vec{p}_2 \rangle &= \frac{-ie_p\mu_{\rm N\Delta}g_{\rm N\Delta\pi}g_{\rm NN\pi}}{(2\pi)^6 36 m^3} \\
\times(\vec{\sigma}_2.\vec{q}_2)\Bigg\{ &\frac{4(m_\Delta - m)\vec{q}_2 + 2|\vec{k}|(i\vec{\sigma}_1 \times \vec{q}_2)}{[(m_\Delta - m)^2 - |\vec{k}|^2](\vec{q}_2^{\,2} + m_\pi^2)}\tau_2^3 \\
&+ \frac{(m_\Delta - m)(i\vec{\sigma}_1 \times \vec{q}_2) + 2|\vec{k}|\vec{q}_2}{[(m_\Delta - m)^2 - |\vec{k}|^2](\vec{q}_2^{\,2} + m_\pi^2)}(i\vec{\tau}_1 \times \vec{\tau}_2)^3 \Bigg\} \times \vec{k} \\
+(1,3 \rightleftharpoons 2,4) & \qquad (3.26)
\end{aligned}
$$

The third and final approximation we consider is obtained by neglecting the $|\vec{k}|$-dependence in the complete static limit description of the baryon propagators. This *soft-photon approximation* gives the conventional description [37] of the $\rm N\Delta\gamma(\pi)$ exchange current for the radiation of a photon of momentum $\vec{k}$,

$$
\begin{aligned}
\langle \vec{p}_3\vec{p}_4 | \vec{J}^{\rm N\Delta}_{\rm eff}(\pi) | \vec{p}_1\vec{p}_2 \rangle &= \frac{-ie_p\mu_{\rm N\Delta}g_{\rm N\Delta\pi}g_{\rm NN\pi}}{(2\pi)^6 36 m^3 (m_\Delta - m)} \\
(\vec{\sigma}_2.\vec{q}_2)&\left\{ \frac{4\tau_2^3\vec{q}_2}{(\vec{q}_2^{\,2} + m_\pi^2)} + \frac{(i\vec{\tau}_1 \times \vec{\tau}_2)^3(i\vec{\sigma}_1 \times \vec{q}_2)}{(\vec{q}_2^{\,2} + m_\pi^2)} \right\} \times \vec{k} \\
+(1,3 \rightleftharpoons 2,4) & \qquad (3.27)
\end{aligned}
$$

We anticipate that this result will differ from the complete static limit whenever the photon energy is comparable to the N-Δ mass difference.

6. *N$\Delta\gamma$ $\rho-$Exchange Currents*

Within the soft-photon approximation, it is well known [37] that the $\rm N\Delta\gamma$ ρ-exchange currents are small compared to the corresponding π-exchange currents, although the destructive interference between the two makes it necessary to include both. In the present work, we anticipate the ρ-exchange contributions to be less important than in other works because the RuhrPot model suggests a very weak $\rm NN\rho$ coupling constant. (In particular, RuhrPot [28] uses $g^2_{\rm NN\rho}/4\pi$=0.2169 and κ_ρ=6.4, whereas (for example) Bonn B [49] adopts $g^2_{\rm NN\rho}/4\pi$=0.92 and κ_ρ=6.1).

Since the $\rm N\Delta\gamma$ ρ-exchange currents are expected to be small, we proceed by taking the *vertex static limit* approximation from the outset. In complete analogy to the derivation of eq. (3.22), we obtain,

$$
\vec{J}^{\rm N\Delta}_{\rm eff}(\rho) = \vec{J}^{\rm N\Delta}_{\rm eff}(\rho;{\rm E}) + \vec{J}^{\rm N\Delta}_{\rm eff}(\rho;{\rm M}) \qquad (3.28)
$$

where,

$$
\begin{aligned}
\langle \vec{p}_3\vec{p}_4 | \vec{J}^{\rm N\Delta}_{\rm eff}(\rho;{\rm E}) | \vec{p}_1\vec{p}_2 \rangle = \big\{ &\big[4N_{\rm E}^{(+)}(\vec{p}_2 + \vec{p}_4) \times \vec{q}_2 + 2N_{\rm E}^{(-)} i\vec{\sigma}_1 \times [(\vec{p}_2 + \vec{p}_4) \times \vec{q}_2]\big]\tau_2^3 \\
&\big[+2N_{\rm E}^{(-)}(\vec{p}_2 + \vec{p}_4) \times \vec{q}_2 + N_{\rm E}^{(+)} i\vec{\sigma}_1 \times [(\vec{p}_2 + \vec{p}_4) \times \vec{q}_2]\big](i\vec{\tau}_1 \times \vec{\tau}_2)^3 \big\} \times \vec{k} + (1,3 \rightleftharpoons 2,4) \qquad (3.29)
\end{aligned}
$$

and

$$
\begin{aligned}
\langle \vec{p}_3\vec{p}_4 | \vec{J}^{\rm N\Delta}_{\rm eff}(\rho;{\rm M}) | \vec{p}_1\vec{p}_2 \rangle = \big\{ &\big[4N_{\rm M}^{(+)}\big[(i\vec{\sigma}_2 \times \vec{q}_2) \times \vec{q}_2\big] - 2N_{\rm M}^{(-)}\big[\vec{\sigma}_1 \times [(\vec{\sigma}_2 \times \vec{q}_2) \times \vec{q}_2]\big]\big]\tau_2^3 \\
&+\big[2N_{\rm M}^{(-)}\big[(i\vec{\sigma}_2 \times \vec{q}_2) \times \vec{q}_2\big] - N_{\rm M}^{(+)}\big[\vec{\sigma}_1 \times [(\vec{\sigma}_2 \times \vec{q}_2) \times \vec{q}_2]\big]\big](i\vec{\tau}_1 \times \vec{\tau}_2)^3 \big\} \times \vec{k} + (1,3 \rightleftharpoons 2,4) \qquad (3.30)
\end{aligned}
$$

with,

$$
\begin{aligned}
N_{\rm M}^{(+)} &= N_0[F^{(1)}_{\rm NN\rho}(Q_2^2) + \kappa_\rho F^{(2)}_{\rm NN\rho}(Q_2^2)][D^\rho_{\rm def} + D^\rho_{\rm abc}], \qquad N_{\rm E}^{(+)} = N_0 F^{(1)}_{\rm NN\rho}(Q_2^2)[D^\rho_{\rm def} + D^\rho_{\rm abc}], \\
N_{\rm M}^{(-)} &= N_0[F^{(1)}_{\rm NN\rho}(Q_2^2) + \kappa_\rho F^{(2)}_{\rm NN\rho}(Q_2^2)][D^\rho_{\rm def} - D^\rho_{\rm abc}], \qquad N_{\rm E}^{(-)} = N_0 F^{(1)}_{\rm NN\rho}(Q_2^2)[D^\rho_{\rm def} - D^\rho_{\rm abc}], \\
N_0 &\overset{\rm SU(6)}{=} \frac{e_p g^2_{\rm NN\rho} G^{\rm V}_{\rm M}(0) G^{\rm N\Delta\rho}_{\rm M}(Q_2^2)}{100 m^3 (2\pi)^6 \omega_\rho(\vec{q}_2)} \qquad (3.31)
\end{aligned}
$$

where we normalize $G^{\rm N\Delta\rho}_{\rm M}(0) = 1 + \kappa_\rho$ and adopt the propagators D^ρ of eq. (3.19). The *complete static limit* and *soft-photon* approximations follow in analogy to the procedures used to develop eqs. (3.26) and (3.27)), the latter resulting in,

$$
\langle \vec{p}_3\vec{p}_4 | \vec{J}^{\rm N\Delta}_{\rm eff}(\rho;{\rm E}) | \vec{p}_1\vec{p}_2 \rangle = \frac{4N_0(1+\kappa_\rho)}{(m_\Delta - m)} \left\{ \frac{4\tau_2^3(\vec{p}_4 + \vec{p}_2) \times \vec{q}_2}{(\vec{q}_2^{\,2} + m_\rho^2)} - \frac{(\vec{\tau}_1 \times \vec{\tau}_2)^3 \vec{\sigma}_1 \times [(\vec{p}_4 + \vec{p}_2) \times \vec{q}_2]}{(\vec{q}_2^{\,2} + m_\rho^2)} \right\} \times \vec{k}
$$

$$
+(1 \rightleftharpoons 2) \qquad (3.32a)
$$

$$
\langle\vec{p}_3\vec{p}_4|\vec{J}^{\,\mathrm{N}\Delta}_{\mathrm{eff}}(\rho;\mathrm{M})|\vec{p}_1\vec{p}_2\rangle = \frac{4N_0(1+\kappa_\rho)^2}{(m_\Delta - m)}\left\{\frac{4\tau_2^3\big[(i\vec{\sigma}_2\times\vec{q}_2)\times\vec{q}_2\big]}{(\vec{q}_2^{\,2}+m_\rho^2)} - \frac{(\vec{\tau}_1\times\vec{\tau}_2)^3\big[\vec{\sigma}_1\times\big[(i\vec{\sigma}_2\times\vec{q}_2)\times\vec{q}_2\big]\big]}{(\vec{q}_2^{\,2}+m_\rho^2)}\right\}\times\vec{k}
$$

$$
+(1 \rightleftharpoons 2) \qquad (3.32b)
$$

where $\vec{J}^{\,\mathrm{N}\Delta}_{\mathrm{eff}}(\rho;\mathrm{M})$ is the conventional result, and $\vec{J}^{\,\mathrm{N}\Delta}_{\mathrm{eff}}(\rho;\mathrm{E})$ is an additional piece (resulting from the convection current part of the NNρ vertex) which is usually ignored on the basis that it is smaller than $\vec{J}^{\,\mathrm{N}\Delta}_{\mathrm{eff}}(\rho;\mathrm{M})$ by a factor of about $1+\kappa_\rho \sim 7$.

C. Direct, Single- and Rescattering Amplitudes

A model-independent expression for the complete invariant amplitude was given in eq. (2.22), where we developed a decomposition into the four terms describing direct, single (*i.e.* initial- and final-state) and rescattering amplitudes shown in Fig. 1. The left-hand side of eq (2.22) is defined by the model-independent expressions of eqs. (2.2) and (2.6), and is Lorentz invariant by construction. However, since the right-hand side of eq. (2.22) is determined by a model-dependent calculation, there can be no *a priori* guarantee that each of the direct, single and rescattering amplitudes are individually Lorentz invariant unless the wave functions are calculated in a manifestly covariant formalism [61].

Since the wave functions are usually constructed from an NN-interaction t-matrix that is defined only in the barycentric frame, and two distinct barycentric frames appear in eq. (2.22), we cast the entire expression into the (maximally symmetric) average barycentric frame, so that the momenta satisfy,

$$
\mathrm{A-frame:} \qquad \vec{p}_1+\vec{p}_2-\tfrac{1}{2}\vec{k} = 0 = \vec{p}_3+\vec{p}_4+\tfrac{1}{2}\vec{k} \qquad (3.33)
$$

and we acknowledge that a formal solution of the initial-, final- and rescattering amplitudes requires the application of boost operators [50–53].

In the following we will make use of the fact that the effective current operator of eq. (3.5) is a totally symmetric under interchange of particles 1 and 2. As such, we obtain in an arbitrary frame,

$$
\begin{aligned}
&\langle\vec{p}_3\vec{p}_4;(s_1s_2)S_f;(t_1t_2)T_f|J_{\mathrm{eff}}|\vec{p}_1\vec{p}_2;(s_1s_2)S_i;(t_1t_2)T_i\rangle\\
&=\langle\vec{p}_4\vec{p}_3;(s_2s_1)S_f;(t_2t_1)T_f|J_{\mathrm{eff}}|\vec{p}_2\vec{p}_1;(s_2s_1)S_i;(t_2t_1)T_i\rangle\\
&=\langle\vec{p}_4\vec{p}_3;(s_1s_2)S_f;(t_1t_2)T_f|J_{\mathrm{eff}}|\vec{p}_2\vec{p}_1;(s_1s_2)S_i;(t_1t_2)T_i\rangle\\
&\quad\times(-1)^{(S_i+S_f+T_i+T_f)} \qquad (3.34)
\end{aligned}
$$

Denoting $|\alpha\rangle = |(s_1s_2)S;(t_1t_2)T\rangle$, we define the anti-symmeterized states as,

$$
|\widetilde{\vec{p}_1\vec{p}_2};\alpha\rangle = \frac{1}{\sqrt{2}}\left\{|\vec{p}_1\vec{p}_2;\alpha\rangle - (-1)^{(S+T)}|\vec{p}_2\vec{p}_1;\alpha\rangle\right\} \qquad (3.35)
$$

It is easy to see that parity conservation is consistent with the Fermi statistics requirement $L+S+T$=odd in the barycentric frame.

1. Direct Amplitudes

The direct amplitudes appearing eq. (2.22a) can be simplified with use of eqs. (3.34) and (3.35) to give,

$$
\begin{aligned}
\mathcal{M}^{\mathrm{D}}_{fi} &= N\vec{\epsilon}(\vec{k},\lambda).\big[\langle\vec{p}_3\vec{p}_4;\alpha_f|\vec{J}_{\mathrm{eff}}|\vec{p}_1\vec{p}_2;\alpha_i\rangle\\
&\quad -(-1)^{(S_i+T_i)}\langle\vec{p}_3\vec{p}_4;\alpha_f|\vec{J}_{\mathrm{eff}}|\vec{p}_2\vec{p}_1;\alpha_i\rangle\big] \qquad (3.36)
\end{aligned}
$$

where N is defined in eq. (2.23) and $\vec{J}_{\mathrm{eff}}$ is given by eq. (3.5). However, momentum conservation demands that the direct terms cannot involve the 1-body part of the effective current density, so in the present numerical results we include the photon coupling to

- NN currents with recoil and wave function renormalization currents.
- N$\bar{\mathrm{N}}$ creation and annihilation currents
- $\rho\pi\gamma$, $\omega\pi\gamma$, $\rho\eta\gamma$ and $\omega\eta\gamma$ exchange currents.
- ΔN currents with π and ρ exchange.

as described by eqs. (3.9), (3.11), (3.16), (3.22), (3.29) and (3.30)

2. Single-Scattering Amplitudes

The initial-state interaction amplitudes appearing eq. (2.22b) are given by,

$$
\begin{aligned}
\mathcal{M}^{\mathrm{I}}_{fi} &= N\vec{\epsilon}(\vec{k},\lambda).\langle\widetilde{\vec{p}_3\vec{p}_4};\alpha_f|\vec{J}_{\mathrm{eff}}(0)]G_i t^{(+)}|\widetilde{\vec{p}_1\vec{p}_2};\alpha_i\rangle\\
&= N\vec{\epsilon}(\vec{k},\lambda).\sum_\alpha\iint d\vec{p}_1^{\,\prime}d\vec{p}_2^{\,\prime}\langle\widetilde{\vec{p}_3\vec{p}_4};\alpha_f|\vec{J}_{\mathrm{eff}}(0)|\vec{p}_1^{\,\prime}\vec{p}_2^{\,\prime};\alpha\rangle\\
&\quad\times\langle\widetilde{\vec{p}_1^{\,\prime}\vec{p}_2^{\,\prime}};\alpha|G_i t^{(+)}|\widetilde{\vec{p}_1\vec{p}_2};\alpha_i\rangle \qquad (3.37)
\end{aligned}
$$

A formal specification of this amplitude follows by inserting the full effective current density of eq. (3.5) and defining boost the operators needed to cast the t-matrix in the initial-state barycentric frame. None of the bremsstrahlung calculations known to us has attempted either of these tasks. Instead, the current has been truncated to include only one-body contributions and boost operators are ignored under the assumption that the part of the invariant amplitude resulting from initial-state correlations alone is itself individually Lorentz invariant.

The first approximation could be removed with a straightforward application of the expressions provided in earlier sections, but since we anticipate the impulse current to be significantly larger than the summed exchange currents, our present numerical applications share the conventional approximation of retaining only the impulse current in the single-scattering amplitudes. The second approximation will be justified in section IV A, where we provide a perturbative analysis that indicates the NNγ impulse-current contributions to the single-scattering terms are close to invariant under Lorentz transformation into the barycentric frames.

We therefore cast the initial-state correlation amplitudes into the initial-state barycentric frame, where

$$\mathrm{I-frame:}\qquad \vec{p}_1+\vec{p}_2=0=\vec{p}_3+\vec{p}_4+\vec{k} \tag{3.38}$$

so that eq. (2.22b) becomes, with the kinematical notation of eq. (3.7),

$$\begin{aligned}\mathcal{M}^{\mathrm{I}}_{fi} &= N\vec{\epsilon}(\vec{k},\lambda).\sum_{\alpha}\langle\alpha_f|\vec{J}_{\mathrm{NN}\gamma}[1](\vec{p}_{3k},\vec{p}_3)|\alpha\rangle\\ &\quad\times\frac{\widetilde{\langle\vec{p}_{3k};\alpha}|t^{(+)}|\widetilde{\vec{p}_1;\alpha_i\rangle}}{2E_1-2E_{3k}+i\eta}\\ &+N\vec{\epsilon}(\vec{k},\lambda).\sum_{\alpha}\langle\alpha_f|\vec{J}_{\mathrm{NN}\gamma}[2](\vec{p}_{4k},\vec{p}_4)|\alpha\rangle\\ &\quad\times\frac{\widetilde{\langle\vec{p}_3;\alpha}|t^{(+)}|\widetilde{\vec{p}_1;\alpha_i\rangle}}{2E_1-2E_3+i\eta}\end{aligned} \tag{3.39}$$

where we have denoted $\langle\vec{p}',-\vec{p}';\alpha_f|t^{(+)}|\vec{p},-\vec{p};\alpha_i\rangle = \langle\vec{p}';\alpha_f|t^{(+)}|\vec{p};\alpha_i\rangle$. In complete analogy, we cast the final-state correlation amplitudes into the final-state barycentric frame, where

$$\mathrm{F-frame:}\qquad \vec{p}_1+\vec{p}_2-\vec{k}=0=\vec{p}_3+\vec{p}_4 \tag{3.40}$$

so that eq. (2.22c) becomes,

$$\begin{aligned}\mathcal{M}^{\mathrm{F}}_{fi} &= N\vec{\epsilon}(\vec{k},\lambda).\sum_{\alpha}\frac{\widetilde{\langle\vec{p}_3;\alpha_f}|t^{(-)\dagger}|\widetilde{\vec{p}_{1k};\alpha\rangle}}{2E_3-2E_{1k}+i\eta}\\ &\quad\times\langle\alpha|\vec{J}_{\mathrm{NN}\gamma}[1](\vec{p}_1,\vec{p}_{1k})|\alpha_i\rangle\\ &+N\vec{\epsilon}(\vec{k},\lambda).\sum_{\alpha}\frac{\widetilde{\langle\vec{p}_3;\alpha_f}|t^{(-)\dagger}|\widetilde{\vec{p}_1;\alpha\rangle}}{2E_3-2E_1+i\eta}\\ &\quad\times\langle\alpha|\vec{J}_{\mathrm{NN}\gamma}[2](\vec{p}_2,\vec{p}_{2k})|\alpha_i\rangle\end{aligned} \tag{3.41}$$

Simple kinematics establishes that the radiation of a real photon implies an off-shell t-matrix, so that we are free to immediately take the limits $\eta\to 0$ in eqs. (3.39) and (3.41).

3. Rescattering Amplitudes

From the results of impulse approximation calculations [9,10,14] we already know that the impulse contributions to the rescattering amplitudes constitute a correction of $\leq 15\%$ to the corresponding single-scattering amplitudes, so for simplicity we neglect from the outset all 2-body currents in the rescattering amplitudes of eq. (2.22d). Hence, in the A-frame of eq. (3.33) we obtain,

$$\begin{aligned}\mathcal{M}^{\mathrm{R}}_{fi} = N\sum_{\alpha\alpha'}\iint d\vec{p}'_1 d\vec{p}'_2\delta^{(3)}(\vec{p}'_1+\vec{p}'_2-\vec{p}_1-\vec{p}_2)\Big\{\\ \langle\vec{p}_3,\widetilde{\vec{p}_4;\alpha_f}|t^{(-)\dagger}G_f|\vec{p}'_{1k},\vec{p}'_2;\alpha_b\rangle\langle\alpha'|J_{\mathrm{NN}\gamma}[1](\vec{p}'_1,\vec{p}'_{1k})|\alpha\rangle\\ \times\langle\vec{p}'_1,\vec{p}'_2;\alpha_a|G_i t^{(+)}|\vec{p}_1,\widetilde{\vec{p}_2;\alpha_i}\rangle\\ +\langle\vec{p}_3,\widetilde{\vec{p}_4;\alpha_f}|t^{(-)\dagger}G_f|\vec{p}'_1,\vec{p}'_{2k};\alpha_b\rangle\langle\alpha'|J_{\mathrm{NN}\gamma}[2](\vec{p}'_2,\vec{p}'_{2k})|\alpha\rangle\\ \times\langle\vec{p}'_1,\vec{p}'_2;\alpha_a|G_i t^{(+)}|\vec{p}_1,\widetilde{\vec{p}_2;\alpha_i}\rangle\Big\}\end{aligned} \tag{3.42}$$

The initial- and final-state barycentric frames differ by the photon momentum, so that no frame can be found where both t-matrices are expressed in their barycentric frame. We therefore introduce a boost operator χ satisfying [50–53],

$$\begin{aligned}|\vec{p}_a,\vec{p}_b\rangle &= \big\{1-i\chi(\vec{\mathcal{P}})\big\}\;|+\vec{p}\rangle|\vec{\mathcal{P}}\rangle+\mathcal{O}(1/m^4),\\ \vec{p} &= \frac{1}{2}(\vec{p}_a-\vec{p}_b),\qquad \vec{\mathcal{P}}=(\vec{p}_a+\vec{p}_b)\end{aligned} \tag{3.43}$$

Eq. (3.42) is manifestly symmetric under interchange of particles 1 and 2. However, for computation purposes, it proves convenient to make use of eqs. (3.34), (3.35) and (3.43) to formally re-express eq. (3.42) in terms of the photon coupling only to nucleon 1,

$$\begin{aligned}\mathcal{M}^{\mathrm{R}}_{fi} &= 2N(-1)^{(S_f+S_i+T_f+T_i)}\sum_{\alpha,\alpha'}\vec{\epsilon}.\int d\vec{p}\langle\vec{p}_3\,\widetilde{+}\,\frac{1}{4}\vec{k};\alpha_f|[1+i\chi(-\frac{\vec{k}}{2})]t^{(-)\dagger}G_f[1-i\chi(-\frac{\vec{k}}{2})]|\vec{p}+\frac{1}{4}\vec{k};\alpha'\rangle\\ &\times\langle\alpha'|\vec{J}_{\mathrm{NN}\gamma}[1](-\vec{p}-\frac{\vec{k}}{2},-\vec{p}+\frac{\vec{k}}{2})|\alpha\rangle\langle+\vec{p}-\frac{1}{4}\vec{k};\alpha|[1+i\chi(+\frac{\vec{k}}{2})]G_i t^{(+)}[1-i\chi(+\frac{\vec{k}}{2})]|\vec{p}_1\,\widetilde{-}\,\frac{1}{4}\vec{k};\alpha_i\rangle\end{aligned} \tag{3.44}$$

where, since the t-matrix conserves spin- and isospin, the sums over the intermediate-state quantum numbers ($\alpha{=}S, M_S, T, M_T$) are restricted such that $S{=}S_i$, $S'{=}S_f$, $T{=}T_i$, $T'{=}T_f$ with conserved isospin projection M_T. For reasons already indicated in the discussion of the single-scattering amplitudes, the boost operators can be neglected in the present work. A recipe for performing the numerical integration over the pole-structures of eq. (3.44) is given in App. C.

IV. RESULTS AND DISCUSSION

A. Impulse Contributions

In the present work we are primarily interested to investigate a consistent calculation of the dominant isoscalar meson-exchange currents in pp-bremsstrahlung. An important precursor to this lies in establishing that the well-known discrepancy between impulse approximation calculations and experimental data cannot be resolved by selecting a different (phase-equivalent) NN-interaction. Although a qualitative similarity exists between the results of recent impulse approximation calculations for pp-bremsstrahlung observables [8–14,16,18] the calculation differences are generally not confined to the differing NN-potentials. Some of these calculations describe the photon coupling to the one-body current via a Foldy-Wouthuysen transformations [8,12,13,16,18] or direct Pauli reduction [10,11,14], whereas others adopt a static limit description [9]. Some include the rescattering amplitudes of Fig. 1c [9–11,14,18] whereas others don't [8,12,13,16]. Further differences are found in the use of relativistic or non-relativistic two-nucleon propagators and/or the application of (guessed) off-shell minimal relativity factors [10,11].

We avoid all of these uncertainties by presenting the results of calculations that are identical apart from the definition of the potential used to generate the t-matrix elements. We also extend the list of commonly compared potentials to include the Bonn [49], Paris [34], Nijmegen [54] and RuhrPot [28]. For the present comparisons we adopt the impulse approximation, so that the effective current density developed in section III B reduces to the sum of (one-body) impulse currents, $\vec{J}_{\rm eff} \sim \vec{J}_{{\rm NN}\gamma}[1] + \vec{J}_{{\rm NN}\gamma}[2]$, as defined in App. B, and is therefore common to all potentials. As such, we retain initial-state, final-state and rescattering amplitudes with the two-nucleon Green's functions described by the relativistic Lippmann-Schwinger propagators. Partial-waves are summed to $J_{\rm max} = 8$ and no form of soft-photon approximation is adopted at any stage.



FIG. 8. Comparison of impulse approximation calculations using Ruhr, Nijmegen, Paris and Bonn potentials and coplanar pp-bremsstrahlung data at $E_{\rm lab}$=280 MeV. The differences between the model results are smaller than their collective discrepancy with experiment.

In Fig. 8 we compare such calculations with the TRIUMF coplanar pp-bremsstrahlung data at $E_{\rm lab}$=280 MeV [35]. The cross section geometries are selected to sample both small and large proton emission angles. The analyzing power geometry is selected on the basis that it is the result most different from zero, and therefore presumably the most reliably measured. Some differences exist between our analyzing power results and those reported elsewhere [11] - primarily due to differences in the rescattering calculation, as discussed in App. C. The essential conclusion here is that impulse approximation calculations using Bonn B, Nijmegen, Paris and RuhrPot wave functions are almost indistinguishable, but exhibit a collective discrepancy with experiment. Given that the purpose of this experiment was to distinguish the predictions of such potentials, the differences between theory and experiment are large.

The final task remaining here is to establish that the impulse current contributions to the single-scattering amplitudes given in eqs (3.39) and (3.41) can be accurately described without boost operators. This approximation is common to all momentum-space bremsstrahlung calculations known to us, yet it appears to have never been verified. Some authors [10,11,20] have sought a solution to the problem by arguing that the non-relativistic t-matrix can be made Lorentz invariant simply by attaching 'minimal relativity' factors [62], so that the Lippmann-Schwinger (LS) equation can be cast in a form that is apparently identical to the Blankenbecler-Sugar (Bbs) equation. However, although both of these integral equations describe the NN interaction in ladder approximation, they are not formally identical because the LS kernel is constructed in a time-ordered (non-covariant) relativistic framework, whereas the BbS-kernel represents one of an infinite number of arbitrary three-dimensional reductions of the covariant Bethe-Salpeter (BS) equation. As such, the Lippmann-Schwinger t-matrix with minimal relativity factors should not be confused with a covariant definition of the NN-interaction [61]. The most serious flaw in the use of minimal relativity factors is, however, that the off-shell factors are completely unknown and must be simply guessed [62]. We regard this procedure as unreasonably arbitrary, particularly since the guessed minimal relativity factors contradict the well known form of the two-nucleon boost operators [50–53].

Since the t-matrix is defined in the conventional way as $t{=}V{+}VGt$ but is available only in the barycentric frame, we consider the arbitrary-frame perturbative reduction of these amplitudes by constructing a toy-model one-boson exchange NN-interaction as,

$$\begin{aligned}&\langle \vec{p}_1^{\,'}, \vec{p}_2^{\,'} | V | \vec{p}_1, \vec{p}_2 \rangle\\ &= \frac{1}{2}\sum_{\beta} \eta H_{\mathrm{NN}\beta}[1](\vec{p}_1, \vec{p}_1^{\,'})\lambda\Big[\frac{1}{\mathcal{E}_i - H_0} + \frac{1}{\mathcal{E}_f - H_0}\Big]\\ &\quad\times \lambda H_{\mathrm{NN}\beta}[2](\vec{p}_2, \vec{p}_2^{\,'})\eta + (1 \rightleftharpoons 2) \qquad (4.1)\end{aligned}$$

where $\beta{=}\pi,\eta,\rho$ and ω, and $H_{\mathrm{NN}\beta}[i]$ is the interaction energy for a meson coupling to nucleon i, as defined in App. B. In Fig. 9 we present the results including only the initial- and final-state interaction amplitudes when the RuhrPot t-matrix is replaced with our toy NN-interaction, the complete expression being cast into the A-, I- and F-frames of eq. (3.33), (3.38), (3.40). We find that the neglect of boost operators represents an error of about 3% or less. We conclude that eq. (3.39) and (3.41) are essentially exact descriptions of the impulse contributions to the single-scattering amplitudes. As such, the application of minimal relativity factors [10,11,20] may need to be reconsidered.



FIG. 9. Initial- and final-state interaction amplitudes in impulse approximation when the t-matrix is replaced with a toy-model boson exchange potential and evaluated in the A-, I-, and F-frames. This justifies the neglect of boost operators in the present work.

B. Exchange Currents and the Relativistic NNγ Vertex

In section III B we showed that a relativistic description of the part of the effective current operator involving the photon coupling to the nucleon current comprises not only the impulse current, but also two-body contributions from the meson-recoil, wave function re-orthonormalization and pair currents. Although relativistic corrections to 1-body NNγ currents have received considerable attention in recent pp-bremsstrahlung works [8,11–13], the 2-body contributions remain to be investigated.

The wave function re-orthonormalization and meson-recoil contributions are not expected to be individually small, but in section III B we recalled [3] that their non-relativistic limits cancel exactly, leaving only purely relativistic effects in the pp-bremsstrahlung observables. In the present numerical applications we retain these contributions in the spirit of exploring the relativistic aspects of the NNγ currents.

By far the most interesting aspect of the relativistic NNγ currents is found in the different manifestations of the N$\bar{\text{N}}$ creation and annihilation pair currents given by various Dyson-equivalent chiral Lagrangians. In particular, the simplest meson-theoretic Lagrangian satisfying PCAC and the chiral commutation relations is the renormalizable σ-model, for which the nucleon-meson interactions are of the form,

$$\mathcal{L} = -g\bar{\psi}[\sigma + i\gamma^5\vec{\tau}.\vec{\pi}']\psi \tag{4.2}$$

With a chiral transformation $\psi \rightarrow (1+\vec{\xi}^2)^{-\frac{1}{2}}\exp[i\gamma^5\vec{\tau}.\vec{\pi}')]\psi$, Weinberg [55] showed that, for a suitably constrained $\vec{\xi}$ and redefined pion fields $\vec{\pi} = (2m/g)\vec{\xi}$, the Lagrangian tranforms to give NNπ and NN$\pi\pi$ interactions with,

$$\mathcal{L}_{\rm pv} = -\frac{g}{2m}\bar{\psi}\left[1+\frac{g^2\vec{\pi}^2}{4m^2}\right]\left[\gamma^5\gamma^\mu\vec{\tau}.\partial_\mu\vec{\pi}+\frac{g}{2m}\vec{\tau}.(\vec{\pi}\times\partial_\mu\vec{\pi})\right]\psi \tag{4.3}$$

Similarly, the chiral transformation $\psi \rightarrow (1+\vec{\xi}^2)^{-\frac{1}{2}}\exp[i(g/2m)\vec{\tau}.(\vec{\pi}'\times\vec{\xi})]\psi$ has been shown by Wess and Zumino [56] to yield NNπ and NN$\pi\pi$ interactions with,

$$\mathcal{L}_{\rm ps} = -\frac{g}{2m}\bar{\psi}\left[1+\frac{g^2\vec{\pi}^2}{4m^2}\right]\left[i\gamma^5\vec{\tau}.\vec{\pi}-\frac{g}{2m}\vec{\pi}^2\right]\psi \tag{4.4}$$

In an elegent summary of these chiral Lagrangians, Gross [57] noted that both $\mathcal{L}_{\rm ps}$ and $\mathcal{L}_{\rm pv}$ give the correct $\pi - N$ scattering lengths and that the NN$\pi\pi$ interactions have ρ and σ quantum numbers.

Our immediate interest lies in the pseudo-vector (pv) and pseudo-scalar (ps) NNπ couplings, both of which are included via the hybrid form of the NNπ Lagrangian we adopt in eq. (3.3) with $0 \leq \lambda \leq 1$. It is a trivial exercise to show that the π-coupling to the positive-frequency components of the nucleon current described in eq. (3.3) is independent of λ, so that non-relativistic calculations cannot differentiate the ps- and pv-couplings. In relativistic applications $\lambda = 0$ is commonly preferred, presumably because the presence of the derivative in the pv-coupling provides for the gauge coupling of a photon to the NNπ vertex, so that the purely isovector non-relativistic seagull exchange currents can be included even when the microscopic structure of the NNπ form factor is ignored. However, there are no formal arguments to rule out $\lambda \neq 0$ and the subject is still under investigation [58,59].

In Fig. 10 we examine the relativistic NNγ currents in pp-bremsstrahlung by comparing the results of calculations with RuhrPot wave functions which neglect (IA) or retain (IA+MEXC) the wave function re-orthonormalization, meson-recoil and N$\bar{\text{N}}$ pair-creation and annihilation currents with purely pv-type (λ=0) and ps-type (λ=1) NNπ couplings.



FIG. 10. RuhrPot results in impulse approximation (IA) as in Fig. 8 compared to corresponding results when the wave function re-orthonormalization, meson-recoil and N$\bar{\text{N}}$-pair currents (IA+MEXC) are included with pseudo-vector (λ=0) or pseudoscalar (λ=1) NNπ interactions. λ can be determined from pp-bremsstrahlung experiments!

The first observation is the most important one: λ can be determined by pp-bremsstrahlung experiments! This is surely the cleanest probe of the Lorentz structure of the NNπ vertex yet considered. We reserve our prediction for λ until we have included the other exchange currents given in section III.

A secondary observation, which we anticipated in ref [14], is that the 2-body contributions are small for $\lambda = 0$. In particular, the λ=0 NNγ MEXC contributions to the cross section at (LEP) θ_3=28.0° and (HEP) θ_4=27.8° are almost entirely negligible. However, the effects are visible in the analyzing powers, as well as in the cross section at θ_3=12.0° and θ_4=12.4°, the latter being reduced at θ_γ=0° and 180° by 0.14 and 0.17 $\mu b/sr^2/rad$ (i.e. 5% and 8%) respectively. While these effects are certainly small, they are already comparable to the model differences shown in Fig. 8.

C. Radiative Vector-Meson Decay Currents

In Fig. 11 we compare the results of calculations using RuhrPot wave functions and the relativistic impulse current which either neglect (IA) or retain (IA+VPγ) the relativistic $\rho\pi\gamma$, $\omega\pi\gamma$, $\rho\eta\gamma$ and $\omega\eta\gamma$ exchange currents [42,43]. The RuhrPot contributions are uniformly small, although we observe a reduction in the cross section at (LEP) θ_3=12.0° and (HEP) θ_4=12.4° at θ_γ=20° by about 0.1 $\mu b/sr^2/rad$ (i.e. 10%).

We have used the fully relativistic expressions of eq (3.16) for the numerical applications of Fig. 11, but to identify the dominant behaviour of these exchange currents it is sufficient to consider only the static limit, neglect the ω- and ρ-meson mass difference and neglect the η-meson contributions altogether. In the non-relativistic limit, this allows us to write,

$$\begin{aligned}\langle\vec{p}_3\vec{p}_4|\vec{J}^{\rho\pi}_{\rm eff}|\vec{p}_1\vec{p}_2\rangle &\overset{\rm nr}{\sim} g_{\rho\pi\gamma}g_{{\rm NN}\rho}[\hat{J}(\vec{q}_1,\vec{q}_2)+\hat{J}(\vec{q}_2,\vec{q}_1)]\vec{\tau}_1.\vec{\tau}_2\\ \langle\vec{p}_3\vec{p}_4|\vec{J}^{\omega\pi}_{\rm eff}|\vec{p}_1\vec{p}_2\rangle &\overset{\rm nr}{\sim} g_{\omega\pi\gamma}g_{{\rm NN}\omega}[\hat{J}(\vec{q}_1,\vec{q}_2)+\hat{J}(\vec{q}_2,\vec{q}_1)]\vec{\tau}^{\,z}_1\end{aligned} \tag{4.5}$$

where

$$\hat{J}(\vec{q}_1,\vec{q}_2) = \frac{ie_p g_{{\rm NN}\pi}(\vec{\sigma}_1.\vec{q}_1)(\vec{q}_1\times\vec{q}_2)}{(2\pi)^6 2m_{\rm V}m[\vec{q}^{\,2}_1+m^2_\pi][\vec{q}^{\,2}_2+m^2_{\rm V}]} \tag{4.6}$$

Since $g_{{\rm NN}\pi}$, $g_{\rho\pi\gamma}$ and $g_{\omega\pi\gamma}$ are essentially fixed by experiment, the freedom in the vector-meson decay exchange currents is limited to the model-dependent values adopted for the experimentally unknown coupling constants $g_{{\rm NN}\rho}$ and $g_{{\rm NN}\omega}$. We have adopted the RuhrPot NN-interaction values of $g_{{\rm NN}\rho}$=1.65 and $g_{{\rm NN}\omega}$=4.95, and note that these values are consistent with the broken SU(3) requirement $g^2_{{\rm NN}\omega}/g^2_{{\rm NN}\rho}$=9. The matrix elements of both isopin operators in eq. (4.5) reduce to unity in pp–bremsstrahlung, so that the $\omega\pi\gamma$ contribution completely dominates the vector-meson decay current contributions and the corresponding $\rho\pi\gamma$ currents are some 15 times smaller.

The magnitude of each PVγ exchange current is, of course, dependent on the choice of NN-interaction. For example, the Bonn B model requires $g_{{\rm NN}\rho}$=3.36 and $g_{{\rm NN}\omega}$=17.5, so that $g^2_{{\rm NN}\omega}/g^2_{{\rm NN}\rho}$=27, which severely contradicts the broken SU(3) prediction of 9. Although the $\rho\pi\gamma$ and $\omega\pi\gamma$ exchange currents have never been included in Bonn model calculations for bremsstrahlung observables, it is easy to see that these currents alone would be respectively more that 4 and 12 times larger than the corresponding results of the present calculation.



FIG. 11. RuhrPot results in impulse approximation (IA) as in Fig. 8 compared to corresponding results when the VPγ= $\rho\pi\gamma + \omega\pi\gamma + \rho\eta\gamma + \omega\eta\gamma$ exchange currents (IA+VPγ) are included. The RuhrPot model has relatively weak NNρ and NNω couplings, so these exchange currents are small, but comparable to the model differences shown in Fig. 8.



FIG. 12. RuhrPot results in impulse approximation (IA) as in Fig. 8 compared to corresponding results with relativistic (rel), vertex-static limit (vsl) and soft-photon approximation (spa) NΔγ π− and ρ−exchange currents. A relativistic description of the NΔγ exchange currents is necessary.



FIG. 13. RuhrPot results in impulse approximation (IA) as in Fig. 8 compared to corresponding results with relativistic NΔγ exchange currents including $\pi + \rho$−exchange [$\pi + \rho$] and π−exchange [π] only. The RuhrPot model has a relatively weak NNρ coupling, so that the ρ−exchange contribution to the NΔγ currents is small.

D. Δ-isobar Currents

In Fig. 12 we compare the results of calculations using RuhrPot wave functions and the relativistic impulse current which either neglect (IA) or retain (IA+Δ) the relativistic NΔγ π− and ρ− exchange currents, as prescribed by the relativistic [rel], vertex-static limit [vsl] and soft-photon approximation [spa] expressions developed in section III B. We do not show results for the complete static limit since these results turn out to be almost indistinguishable from the vertex static limit. In Fig. 13 we decompose the contributions to the relativistic NΔγ currents into π− and ρ− exchange contributions. Collectively, these figures show that the RuhrPot description of the NΔγ exchange currents are very large and require



FIG. 14. Contributions from individual currents to the cross section. These are (IA) = impulse current including initial-, final- and rescattering correlations, (N$\Delta\gamma$) with summed $\pi-$ and $\rho-$exchange, (VPγ) = $\rho\pi\gamma$ + $\omega\pi\gamma$ + $\rho\eta\gamma$ + $\omega\eta\gamma$, (WFRR+Pair) = NNγ wave function re-orthonormalization and meson recoil and $\bar{\text{N}}$N-pair creation and annihilation currents with $\lambda = 0$. (Full) denotes the sum of all these currents (with interferences).

a relativistic description, but that the $\rho-$ exchange contribution is comparatively small.

E. Comparison of Exchange Currents

In Fig. 14 we examine the individual contributions of each of the 1- and 2-body currents developed in the previous sections with the complete effective current of the present work. We observe that the N$\Delta\gamma$ exchange currents are substantially larger than all other contributions and we recall from Fig. 13 that, in the RuhrPot description, these are completely dominated by the $\pi-$exchange contributions. We observe minor but non-negligible contributions from the VPγ currents, and we recall from sec IV C that the magnitude of these contributions is essentially determined by the NNω coupling constant in the NN interaction.

F. Comparison With Experiment

In Figs. 15-18 we compare results of calculations using RuhrPot wave functions and the relativistic impulse current which either neglect (IA) or retain (IA+MEXC) all of the relativistic meson-exchange currents developed in the present work. These include the wave function re-orthonormalization and meson-recoil currents, the N$\bar{\text{N}}$ creation and annihilation pair-currents (for both ($\lambda = 0$) pv- and ($\lambda = 1$) ps- NNπ-couplings), the $\rho\pi\gamma$, $\omega\pi\gamma$, $\rho\eta\gamma$ and $\omega\eta\gamma$ exchange currents and the N$\Delta\gamma$ currents with $\pi-$ and $\rho-$ exchange contributions. We observe that the sensitivity to the admixture of ps ($\lambda = 1$) and pv ($\lambda = 0$) couplings survives when the currents are consistently combined.

The inclusion of the relativistic exchange currents can be seen to provide a reasonable description of the cross section data in Figs. 15-16, but Figs. 17-18 show that a persistent discrepancy with experiment remains for small θ_3 and large θ_4. The impulse approximation results for the cross section data give χ^2_{ia}/datum = 5.8, whereas the complete exchange current calculations with ps and pv NNπ couplings giving χ^2_{ps}/datum = 6.1 and χ^2_{pv}/datum = 4.7 respectively. Moreover, Fig. 15-16 shows that adopting a ps NNπ coupling produces structure in the cross section that is simply absent in the data. As such, the present calculations indicate that the data favours $\lambda \sim 0$, although Figs. 17-18 show that some of the most serious discrepancies with experiment cannot be resolved in terms of the NNπ coupling alone.

The importance of our conclusions on the Lorentz structure of the NNπ vertex are contingent on a reliable data set, so we vigorously stress the need for more precise measurements of all observables where the ps- and pv-couplings give very different results.

G. Some problems with non-perturbative descriptions

Strong interaction transitions between NN, ΔN and $\Delta\Delta$ states can be described non-perturbatively with a coupled channel t-matrix [31–33] and indeed have already been used to calculate pp-bremsstrahlung observables [19,20]. Under these circumstances it may appear curious that we have chosen to present a perturbative description of such amplitudes as meson-exchange currents. There are, however, a number of difficulties in applications of these coupled channel t-matrices, the most serious of which appears to arise from the inconsistencies that exist between the Paris [34] NN$\rightleftharpoons$NN and the Ried parameterized version of the static-limit Bochum [31–33] ΔN $\rightleftharpoons$NN interaction. In particular, the conflicting definitions of the NNπ and N$\Delta\pi$ coupling constants and form factors makes it impossible to reliably remove the



FIG. 15. Coplanar pp-bremsstrahlung data at E_{lab}=280 MeV and $\theta_4 = 12.4°$ compared to RuhrPot calculations including (IA+MEXC) or excluding (IA) relativistic meson-exchange currents with pv (λ=0) or ps (λ=1) NNπ interactions. The exchange currents include wave function re-orthonormalization and meson-recoil currents, N$\bar{\mathrm{N}}$ pair creation and annihilation currents, $\rho\pi\gamma$ + $\omega\pi\gamma$ + $\rho\eta\gamma$ + $\omega\eta\gamma$ vector-meson decay currents and N$\Delta\gamma(\pi, \rho, \Gamma_\Delta$=115 MeV) exchange currents.



FIG. 16. Same as Fig. 15 except that $\theta_4 = 17.3°$.



FIG. 17. Same as Fig. 15 except that $\theta_4 = 21.2°$.



FIG. 18. Same as Fig. 15 except that $\theta_4 = 27.8°$.

double-counted NN⇌NN amplitudes involving intermediate NΔ states, so that a free parameter is introduced to simulate the necessary subtraction. Any specification of the two-body currents would suffer a similar ambiguity. Finally, the assumed Lorentz invariance of the NN ⇌ ΔN transition t-matrices remains to be investigated.

This leads us to consider a generalization of our formalism to obtain a fully consistent and microscopic description of the non-perturbative transitions between the NN, ΔN and ΔΔ states. More precisely, we will identify two minimum requirements of such an approach that appear to have been been neglected to date.

Recalling from sec. II B the freedom to choose any desired partition of the total Hilbert space, we now modify our earlier choice so that the Δ degrees of freedom are included in the η-space. (Details can be found in App A). Within this approach, the leading order contributions involve not only the NNγ impulse, wave function re-orthonormalization and meson-recoil terms, but also NΔγ initial- and final-state correlations terms and NΔγ wave function re-orthonormalization and meson-recoil terms, as shown in Fig. 19. Our earlier specification of the NNγ one- and two-body currents remains unchanged and will not be further discussed here. The additional leading-order contributions involving Δ isobars are given by,

$$\begin{aligned}\langle \vec{p}_3\vec{p}_4|\vec{J}^{\mathrm{N}\Delta}_{11}|\vec{p}_1\vec{p}_2\rangle = -g_{\sigma\tau}\sum_{\beta}\Big[&\\ D^{\beta}_1 H^{\sigma}_{\mathrm{N}\Delta\beta}[1](\vec{p}_{1k},\vec{p}_3)\vec{J}_{\Delta\mathrm{N}\gamma}[1](\vec{p}_1,\vec{p}_{1k})H^{\tau}_{\mathrm{NN}\beta}[2](\vec{p}_2,\vec{p}_4)&\\ +D^{\beta}_3\vec{J}_{\mathrm{N}\Delta\gamma}[1](\vec{p}_{3k},\vec{p}_3)H^{\sigma}_{\Delta\mathrm{N}\beta}[1](\vec{p}_1,\vec{p}_{3k})H^{\tau}_{\mathrm{NN}\beta}[2](\vec{p}_2,\vec{p}_4)\Big]&\\ +(1,3\rightleftharpoons 2,4)&\end{aligned}\qquad(4.7)$$

where $\beta=\vec{\pi}$ or $\vec{\rho}$ and the factor $-g_{\sigma\tau}$ and all references to the Lorentz indices σ and τ are to be ignored for the scalar mesons. The exact form of D^{β} is given in app A, but here it is sufficient to note the non-relativistic limit,

$$D^{\beta}_1 \overset{\mathrm{nr}}{\sim} D^{\beta}_3 \overset{\mathrm{nr}}{\sim} \frac{-1}{2\omega_{\beta}(\vec{q}_2)(m_{\Delta}-m+\omega_{\beta})} \times\left[1+\frac{(m_{\Delta}-m)^2}{\omega_{\beta}(\vec{q}_2)[m_{\Delta}-m+\omega_{\beta}(\vec{q}_2)]}\right]\qquad(4.8)$$

The first term in the brackets of eq (4.8) represents the NΔγ initial- and final-state strong-interaction amplitudes, and the second term gives the NΔγ wave function re-orthonormalization and meson-recoil currents. Unlike their analogous NNγ terms, where only one-body currents survive in the static limit, the NΔγ wave function re-orthonormalization and meson-recoil currents do *not* vanish in static limit - even for soft photons (cf. ref [20]). Indeed noting that $m_{\Delta}-m\sim 2m_{\pi}$ shows that these contributions make an important contribution to the intermediate-state ΔN amplitudes in the low energy observables.



FIG. 19. NΔγ wave function re-orthonormalization and meson-recoil exchange currents. These currents are necessary to preserve the orthonormality of the initial- and final-state wave functions described by NN, ΔN coupled channel transition t-matrices.

Since the coupled channel t-matrices have been calculated only in the barycentric frame, it remains to either calculate their boost operators, or demonstrate that they are effectively Lorentz invariant. In sec IV A (see Fig. 9) we demonstrated such invariance for a toy-model boson-exchange potential and commented on the reliability of guessing minimal relativity factors. We anticipate that a similar approximate invariance will probably hold for the coupled channel NN⇌NN t-matrices involving intermediate ΔN-states. However, in Fig. 20 we observe that the corresponding NΔγ initial- and final-state interaction amplitudes calculated in the average barycentric frame of eq. (3.33) are seriously different from the corresponding results evaluated in the I- and F-frames of eqs. (3.38) and (3.40). This shows that the leading-order contributions to the coupled channel ΔN ⇌NN and NN⇌ ΔN t-matrices are poorly approximated under the assumption that they are Lorentz invariant.

We conclude that a non-perturbative description of the Δ isobar amplitudes via coupled channel t-matrix calculations involves two complications:



FIG. 20. Comparison of perturbative NΔ initial- and final-state correlation amplitudes calculated in the (average barycentric) A-frame and compared to corresponding results that are obtained in the initial- and final-state barycentric frames. The discrepancies show the need for boost operators and indicate errors of about 20% result when the NN$\rightleftharpoons$ ΔN t-matrix is assumed to be Lorentz invariant.

- There are non-vanishing contributions from the N$\Delta\gamma$ wave function re-orthonormalization and meson-recoil terms. The neglect of these two-body currents contributes serious errors at low energies.

- The initial- and final-state N$\Delta\gamma$ interactions are poorly approximated in the absence of boost operators. The neglect of boost operators contributes serious errors at high energies.

As such, a meaningful specification of non-perturbative Δ contributions requires the calculation of two-body currents and boost operators. This has not been recognized in the past. Although the first requirement can easily be satisfied by retaining a subset of the exchange currents we have presented in this work, there are outstanding problems that need to be solved if boost operators are to be defined beyond $\mathcal{O}(1/m^4)$. Such developments stand as a challenge for future theoretical work but can only be approached within a model providing a consistent and microscopic description of all meson-baryon dynamics

H. Relativistic Effects

In earlier work [1] we presented selected pp-bremsstrahlung observables calculated using RuhrPot wave functions and the relativistic impulse current. The re-scattering contributions of Fig. 1c were retained, as were the relativistic $\rho\pi\gamma$, $\omega\pi\gamma$, $\rho\eta\gamma$ and $\omega\eta\gamma$ exchange currents. We also included the N$\Delta\gamma$ currents with $\pi-$ and $\rho-$ exchange in the complete static limit, as defined in sec III B 5. No form of soft-photon approximation was adopted at any stage.

In the present work we have extend the exchange currents to include the wave function re-orthonormalization and meson-recoil currents of eq. (3.9) and the N$\bar{\text{N}}$-pair creation and annihilation currents (for both ($\lambda = 0$) pv- and ($\lambda = 1$) ps- NNπ-couplings) of eq. (3.11) that are required for a truly relativistic description of the NNγ vertex. We have also replaced the complete static limit description of the dominant N$\Delta\gamma(\pi)$ exchange current with the relativistic up-grade of eq. (3.22).

In Fig. 21 we consider the magnitude of these purely relativistic effects by comparing with our earlier descriptions of the meson-exchange currents. We consider here only the pv NNπ coupling ($\lambda = 0$). From Figs 10,12 and 13 we already know that the largest difference between these exchange current results stems from the relativistic corrections to the N$\Delta\gamma(\pi)$ contribution. In addition, from Fig. 12 and sections III B 5 and IV D we note that this difference results from the neglect of the N$-\Delta$ mass difference in the complete static limit exchange current operators used in ref. [1]. Although we find no need to change the qualitative conclusions reported in ref [1], a comparison of pp-bremsstrahlung calculations with experimental data near the π-production threshold clearly requires a relativistic description of the isoscalar meson-exchange currents.

V. CONCLUSIONS

We have presented a parameter-free and relativistic extension of the RuhrPot meson-baryon model to define the dominant isoscalar meson-exchange currents. These include the first relativistic calculations for the wave function re-orthonormalization, meson-recoil and N$\bar{\text{N}}$-pair creation and annihilation currents in pp-bremsstrahlung. We also included the fully relativistic $\rho\pi\gamma$, $\omega\pi\gamma$, $\rho\eta\gamma$ and $\omega\eta\gamma$ currents and a relativistic upgrade to our earlier N$\Delta\gamma(\pi,\rho)$ exchange currents [1,15].

The results of these calculations show that the meson-exchange contributions to the pp-bremsstrahlung observables below the $\pi-$production threshold are large. As



FIG. 21. Selected results from figs 15-18 using the ps NNπ coupling with (Full) and without (IA) the relativistic exchange currents, now compared with the (CSL) results of ref [1]. The CSL results differ from the full ones in that the wave function re-orthonormalization, meson recoil and N$\bar{\text{N}}$-pair currents are neglected entirely and the N$\Delta\gamma(\pi)$ exchange current is taken in the complete static limit. The largest numerical difference in the meson-exchange current results can be traced to the neglect of the N$-\Delta$ mass difference in the CSL operators. See also sections III B 5 and IV D for a detailed discussion.

such, a meaningful interpretation of experiment obviously requires a completely consistent description of the meson-baryon dynamics defining the NN-interaction, the exchange currents and the form factors that they contain.

Although the wave function re-normalization and meson-recoil contributions are well known to cancel in the static limit for soft photons, this cancellation is poorly satisfied in bremsstrahlung experiments where the kinematics has been contrived to maximize the photon energy. As such, retaining these contributions is necessary if the orthonormality of the wave functions is to be preserved. Although these two-body currents have never before been included in bremsstrahlung calculations, they are necessary for a relativistic description of the NNγ vertex.

The motivation for developing such a relativistic scheme is found in one of the oldest outstanding puzzles in nuclear physics. The Lorentz structure of the NNπ Lagrangian $\mathcal{L}_{\text{NN}\pi}$ is universally accepted to comprise an unknown mixture of ps ($\lambda = 1$) and pv ($\lambda = 0$) couplings. This mixture cannot be distinguished by any non-relativistic calculation and, for quite different reasons, makes almost no impact on relativistic calculations for the $N\pi$ scattering lengths. However, within our relativistic framework we have shown that the existing pp-bremsstrahlung data indicates λ is small. This is surely the most reliable assessment of the NNπ Lagrangian available to date.

We quantified the VPγ exchange current contributions and noted their obvious relationship to the NN-interaction. While the small vector-meson couplings in the RuhrPot NN interaction render these effects no larger than the NN-interaction model differences in impulse approximation, we noted the necessity to include these currents in calculations for models using large NNω and NNρ couplings (e.g. Bonn B). However, the largest isovector exchange current in pp-bremsstrahlung results from intermediate-state isobar excitation via π-exchange. We investigated a series of approximations for this current and found that a relativistic description is necessary. Given the practical importance of this result, a compact closed form expression was provided and the sources of error in various approximations were identified.

We demonstrated our earlier assertion [1,15] that the N$\Delta\gamma$ wave function re-orthonormalization and meson-recoil contributions do *not* vanish - even in the static limit for soft photons. Their neglect in recent applications [19,20] indicates that the the orthonormality of the initial and final state wave functions is not preserved. In a perturbative analysis we showed that the assumed Lorentz invariance of the NN$\rightleftharpoons$NN interactions is accurate to about 3%, but that a similar assumption [19,20] for the NN$\rightleftharpoons$ ΔN interaction implies unacceptable errors of around 20%. We noted that a non-perturbative development of our parameter-free calculations requires inclusion of the N$\Delta\gamma$ wave function re-orthonormalization and meson-recoil exchange currents (for which exact expressions were provided) and the application of boost operators. The need for more precise experimental data has been stressed.

ACKNOWLEDGMENTS

This work is supported by COSY-KFA Jülich (41140512), the Deutsche Forschungsgemeinschaft (Ga 153/11-4) and BMFT.

APPENDIX A: NON-PERTURBATIVE N$\Delta\gamma$ AMPLITUDES

Within the formalism of sec II B, a non-perturbative description of the Δ isobar contributions requires our partition of the Hilbert space to modified such that,

$$\mathcal{H}_{\eta_1} = \Big\{|\mathrm{NN}\rangle\Big\}, \quad \mathcal{H}_{\eta_2} = \Big\{|\Delta\mathrm{N}\rangle\Big\}, \quad \mathcal{H}_{\eta_3} = \Big\{|\mathrm{N}\Delta\rangle\Big\},$$
$$\mathcal{H}_{\eta_4} = \Big\{|\Delta\Delta\rangle\Big\}, \quad \mathcal{H}_{\lambda} = \Big\{|\text{the rest}\rangle\Big\} \qquad \text{(A1)}$$

with projection operators satisfying $\eta = \sum_{i=1}^{4} \eta_i$ and $\eta_i \eta_j = \eta_i \delta_{ij}$. Denoting an arbitrary operator causing transitions from the η_i-space to the η_j-space as $\eta_j \mathcal{O} \eta_i = \mathcal{O}_{ji}$, we require (in analogy to eq (2.22)) matrix elements of the form,

$$\begin{aligned}[\mathcal{M}_{fi}]_{11} = N\vec{\epsilon}(\vec{k},\lambda) \sum_{i,j=1}^{4} \langle \vec{p}_3 \widetilde{\vec{p}_4; \alpha}_f | \eta_1 [\mathbb{1} + t_{1i}^{(-)\dagger} G_{1i}] \\ \times [\vec{J}_{\mathrm{eff}}]_{ij} [\mathbb{1} + G_{j1} t_{j1}^{(+)}] \eta_1 | \vec{p}_1 \widetilde{\vec{p}_2; \alpha}_i \rangle \qquad \text{(A2)}\end{aligned}$$

where we have denoted the effective 1+2-body current density as

$$\begin{aligned}[\vec{J}_{\mathrm{eff}}]_{ij} = \eta_i \Big[J + JA + A^\dagger J + A^\dagger JA \\ - \frac{1}{2} J A^\dagger A - \frac{1}{2} A^\dagger A J + \cdots \Big] \eta_j \qquad \text{(A3)}\end{aligned}$$

First consider the amplitudes involving an effective current describing NN$\rightarrow$NN transitions, which must therefore be taken between inital- and final-state wave functions constructed from the NN$\rightarrow$NN t-matrix. These amplitudes include the dominant NNγ impulse currents, as well as contributions from the NNγ wave function re-orthonormalization and meson-recoil currents. (Adopting (A1) does not alter the exact expressions for these currents given by eq. (3.9) and (3.10)). However, we now have additional contributions, as shown in Fig. 19, which are given by eq. (4.7) with

$$\begin{aligned}D_1^\beta = & \frac{1}{[E_3 - E_{\Delta 1k} - \omega_\beta(\vec{q}_2)][E_2 - E_4 - \omega_\beta(\vec{q}_2)]} \\ & - \frac{1}{2[E_4 - E_2 - \omega_\beta(\vec{q}_2)][E_1 - E_3 - \omega_\beta(\vec{q}_2)]} \\ & - \frac{1}{2[E_3 - E_{\Delta 1k} - \omega_\beta(\vec{q}_2)]} \\ & \quad \times \frac{1}{[E_1 + E_2 - E_{\Delta 1k} - E_4 - \omega_\beta(\vec{q}_2)]} \\ D_3^\beta = & \frac{1}{[E_4 - E_2 - \omega_\beta(\vec{q}_2)][E_1 - E_{\Delta 3k} - \omega_\beta(\vec{q}_2)]} \\ & - \frac{1}{2[E_3 - E_1 - \omega_\beta(\vec{q}_2)][E_2 - E_4 - \omega_\beta(\vec{q}_2)]} \\ & - \frac{1}{2[E_3 + E_4 - E_{\Delta 3k} - E_2 - \omega_\beta(\vec{q}_2)]} \\ & \quad \times \frac{1}{[E_1 - E_{\Delta 3k} - \omega_\beta(\vec{q}_2)]} \qquad \text{(A4)}\end{aligned}$$

where $\beta = \vec{\pi}$ or $\vec{\rho}$. Unlike their analogous NNγ terms, these N$\Delta\gamma$ wave function re-orthonormalization and meson-recoil currents do *not* vanish in the static limit. Their inclusion is required to preserve the orthonormality of the hadronic wave functions.

Next consider the amplitudes involving an effective current describing ΔN $\rightleftharpoons$NN transitions, which must therefore be taken between inital- and final-state wave functions constructed from ΔN ⇌NN and NN→NN t-matrices.

$$
\begin{aligned}
[\mathcal{M}_{fi}]_{11} = {} & N\vec{\epsilon}(\vec{k},\lambda).\langle\widetilde{\vec{p}_3\vec{p}_4;\alpha_f}|\eta_1[\mathbb{1}+t_{11}^{(-)\dagger}G_{11}]\\
& \times[\vec{J}_{\rm eff}]_{12}[G_{21}t_{21}^{(+)}]\eta_1|\widetilde{\vec{p}_1\vec{p}_2;\alpha_i}\rangle\\
& + N\vec{\epsilon}(\vec{k},\lambda).\langle\widetilde{\vec{p}_3\vec{p}_4;\alpha_f}|\eta_1[t_{12}^{(-)\dagger}G_{12}]\\
& \times[\vec{J}_{\rm eff}]_{21}[\mathbb{1}+G_{11}t_{11}^{(+)}]\eta_1|\widetilde{\vec{p}_1\vec{p}_2;\alpha_i}\rangle\\
& + (1,3\rightleftharpoons 2,3) \qquad\qquad (\mathrm{A5})
\end{aligned}
$$

These are the N$\Delta\gamma$ initial- and final-state interaction amplitudes and should obviously be specified in a consistent frame. However, the t-matrices are available only in the barycentric frame, yet are required for the initial- and final-state interactions of eq. (A5) in barycentric frames which differ by the photon momentum. The procedure adopted in recent works [19,20] is to attach (guessed) off-shell minimal relativity factors and assume this renders the initial- and final-state interaction terms individually Lorentz invariant. Rather than adopt this assumption, we provide an exact specification of the leading-order contributions to these amplitudes via eq. (4.7) with,

$$
\begin{aligned}
D_1^\beta = {} & \frac{1}{2(E_3+E_4-E_{\Delta 1k}-E_2)}\\
& \times\Big[\frac{1}{(E_3-E_{\Delta 1k}-\omega_\beta)}-\frac{1}{(E_4-E_2-\omega_\beta)}\Big]\\
D_3^\beta = {} & \frac{1}{2(E_1+E_2-E_{\Delta 3k}-E_4})\\
& \times\Big[\frac{1}{(E_1-E_{\Delta 3k}-\omega_\beta)}-\frac{1}{(E_2-E_4-\omega_\beta})\Big] \qquad (\mathrm{A6})
\end{aligned}
$$

APPENDIX B: VERTEX FUNCTIONS

1. Interaction energies for strong vertices

For a meson with momentum $\vec{q}$ and mass m_β we define,

$$
N_\beta = g_{\mathrm{NN}\beta}\left\{\frac{\mathcal{E}_i\mathcal{E}_f}{(2\pi)^3 8\omega_\beta E_f E_i}\right\}^{1/2}, \qquad \omega_\beta=\sqrt{(\vec{q})^2+m_\beta^2} \tag{B1}
$$

The interaction energies $\langle 0|b(\vec{p}_f)\; -\int d^3x\Gamma_{\mathrm{NN}\beta}\, b^\dagger(\vec{p}_i)|0\rangle$ for the coupling of mesons to the positive frequency nucleon current are,

$$
H_{\mathrm{NN}\pi}=N_\pi h_{\mathrm{NN}\pi}\,\vec{\tau}, \qquad H_{\mathrm{NN}\rho}=N_\rho\epsilon_\mu h^\mu_{\mathrm{NN}\rho}\,\vec{\tau}, \qquad H_{\mathrm{NN}\delta}=N_\delta h_{\mathrm{NN}\delta}\,\vec{\tau}, \tag{B2a}
$$

$$
h_{\mathrm{NN}\pi}=-iF_{\mathrm{NN}\pi}\vec{\sigma}.\left[\frac{\vec{p}_f}{\mathcal{E}_f}-\frac{\vec{p}_i}{\mathcal{E}_i}\right] \tag{B2b}
$$

$$
h^0_{\mathrm{NN}\rho}=\left[F^{(1)}_{\mathrm{NN}\rho}+\kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\Big[1-\frac{E_f+E_i}{2m}\Big]\right]+\left[F^{(1)}_{\mathrm{NN}\rho}+\kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\Big[1+\frac{E_f+E_i}{2m}\Big]\right]\left[\frac{\vec{p}_f.\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i}+\frac{i\vec{\sigma}.(\vec{p}_f\times\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i}\right] \tag{B2c}
$$

$$
\vec{h}_{\mathrm{NN}\rho}=\left[F^{(1)}_{\mathrm{NN}\rho}+\kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\right]\left[\frac{\vec{p}_f}{\mathcal{E}_f}+\frac{\vec{p}_i}{\mathcal{E}_i}+i\vec{\sigma}\times\left[\frac{\vec{p}_f}{\mathcal{E}_f}-\frac{\vec{p}_i}{\mathcal{E}_i}\right]\right]-\frac{\kappa_\rho F^{(2)}_{\mathrm{NN}\rho}}{2m}(\vec{p}_f+\vec{p}_i)\left[1-\frac{\vec{p}_f.\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i}-\frac{i\vec{\sigma}.(\vec{p}_f\times\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i}\right] \tag{B2d}
$$

$$
h_{\mathrm{NN}\delta}=F_{\mathrm{NN}\delta}\left[1-\frac{\vec{p}_f.\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i}+\frac{i\vec{\sigma}.(\vec{p}_f\times\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i}\right] \tag{B2e}
$$

The interaction energies $\langle 0|b(\vec{p}_f)\; -\int d^3x\Gamma_{\mathrm{NN}\beta}\, d(-\vec{p}_i)|0\rangle$ for meson couplings to the pair-creation current are,

$$
H_{\bar{\mathrm{N}}\mathrm{N}\pi}=N_\pi h_{\bar{\mathrm{N}}\mathrm{N}\pi}\vec{\tau}, \qquad H_{\bar{\mathrm{N}}\mathrm{N}\rho}=N_\rho\epsilon_\mu h^\mu_{\bar{\mathrm{N}}\mathrm{N}\rho}\,\vec{\tau}, \qquad H_{\bar{\mathrm{N}}\mathrm{N}\delta}=N_\delta h_{\bar{\mathrm{N}}\mathrm{N}\delta}\,\vec{\tau} \tag{B3a}
$$

$$
h_{\bar{\mathrm{N}}\mathrm{N}\pi}=+iF_{\mathrm{NN}\pi}\left\{\Big[1-(1-\lambda)\frac{E_i}{m}\Big]+\Big[1+(1-\lambda)\frac{E_i}{m}\Big]\Big[\frac{\vec{p}_f.\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i}+i\vec{\sigma}.\Big(\frac{\vec{p}_f\times\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i}\Big)\Big]\right\} \tag{B3b}
$$

$$
h^0_{\bar{\mathrm{N}}\mathrm{N}\rho}=\left[F^{(1)}_{\mathrm{NN}\rho}+\kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\Big[1+\frac{E_f-E_i}{2m}\Big]\right]\frac{\vec{\sigma}.\vec{p}_f}{\mathcal{E}_f}-\left[F^{(1)}_{\mathrm{NN}\rho}+\kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\Big[1-\frac{E_f-E_i}{2m}\Big]\right]\frac{\vec{\sigma}.\vec{p}_i}{\mathcal{E}_i} \tag{B3c}
$$

$$
\begin{aligned}
\vec{h}_{\bar{\mathrm{N}}\mathrm{N}\rho}= {} & \left[F^{(1)}_{\mathrm{NN}\rho}+\kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\Big[1-\frac{E_i}{m}\Big]\right]\vec{\sigma}+\frac{\kappa_\rho F^{(2)}_{\mathrm{NN}\rho}}{2m}(\vec{p}_f+\vec{p}_i)\left[\frac{\vec{\sigma}.\vec{p}_f}{\mathcal{E}_f}+\frac{\vec{\sigma}.\vec{p}_i}{\mathcal{E}_i}\right]\\
& +\left[F^{(1)}_{\mathrm{NN}\rho}+\kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\Big[1+\frac{E_i}{m}\Big]\right]\left[\frac{i(\vec{p}_f\times\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i}+\frac{\vec{\sigma}(\vec{p}_f.\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i}-\frac{\vec{p}_f(\vec{\sigma}.\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i}-\frac{(\vec{\sigma}.\vec{p}_f)\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i}\right]
\end{aligned} \tag{B3d}
$$

$$h_{\bar{\mathrm{N}}\mathrm{N}\delta} = -F_{\mathrm{NN}\delta}\left[\frac{\vec{\sigma}.\vec{p}_f}{\mathcal{E}_f} + \frac{\vec{\sigma}.\vec{p}_i}{\mathcal{E}_i}\right] \tag{B3e}$$

The interaction energies $\langle 0|d^\dagger(-\vec{p}_f) - \int d^x \Gamma_{\mathrm{NN}\beta}\, b^\dagger(\vec{p}_i)|0\rangle$ for meson couplings to the pair-annihilation current are,

$$H_{\mathrm{N}\bar{\mathrm{N}}\pi} = N_\pi h_{\mathrm{N}\bar{\mathrm{N}}\pi}\,\vec{\tau}, \qquad H_{\mathrm{N}\bar{\mathrm{N}}\rho} = N_\rho \epsilon_\mu h^\mu_{\mathrm{N}\bar{\mathrm{N}}\rho}\,\vec{\tau} \qquad H_{\mathrm{N}\bar{\mathrm{N}}\delta} = N_\delta h_{\mathrm{N}\bar{\mathrm{N}}\delta}\,\vec{\tau} \tag{B4a}$$

$$h_{\mathrm{N}\bar{\mathrm{N}}\pi} = -iF_{\mathrm{NN}\pi}\Bigg\{\Big[1-(1-\lambda)\frac{E_f}{m}\Big] + \Big[1+(1-\lambda)\frac{E_f}{m}\Big]\Big[\frac{\vec{p}_f.\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i} + i\vec{\sigma}.\Big(\frac{\vec{p}_f\times\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i}\Big)\Big]\Bigg\} \tag{B4b}$$

$$h^0_{\mathrm{N}\bar{\mathrm{N}}\rho} = \left[F^{(1)}_{\mathrm{NN}\rho} + \kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\Big[1-\frac{E_f-E_i}{2m}\Big]\right]\frac{\vec{\sigma}.\vec{p}_i}{\mathcal{E}_i} - \left[F^{(1)}_{\mathrm{NN}\rho} + \kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\Big[1+\frac{E_f-E_i}{2m}\Big]\right]\frac{\vec{\sigma}.\vec{p}_f}{\mathcal{E}_f} \tag{B4c}$$

$$\begin{aligned}\vec{h}_{\mathrm{N}\bar{\mathrm{N}}\rho} = {}& \left[F^{(1)}_{\mathrm{NN}\rho} + \kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\Big[1-\frac{E_f}{m}\Big]\right]\vec{\sigma} + \frac{\kappa_\rho F^{(2)}_{\mathrm{NN}\rho}}{2m}(\vec{p}_f+\vec{p}_i)\left[\frac{\vec{\sigma}.\vec{p}_f}{\mathcal{E}_f} + \frac{\vec{\sigma}.\vec{p}_i}{\mathcal{E}_i}\right] \\ &+ \left[F^{(1)}_{\mathrm{NN}\rho} + \kappa_\rho F^{(2)}_{\mathrm{NN}\rho}\Big[1+\frac{E_f}{m}\Big]\right]\left[\frac{i(\vec{p}_f\times\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i} + \frac{\vec{\sigma}(\vec{p}_f.\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i} - \frac{\vec{p}_f(\vec{\sigma}.\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i} - \frac{(\vec{\sigma}.\vec{p}_f)\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i}\right]\end{aligned} \tag{B4d}$$

$$h_{\bar{\mathrm{N}}\mathrm{N}\delta} = -F_{\mathrm{NN}\delta}\left[\frac{\vec{\sigma}.\vec{p}_f}{\mathcal{E}_f} + \frac{\vec{\sigma}.\vec{p}_i}{\mathcal{E}_i}\right] \tag{B4e}$$

where all strong form factors are evaluated at $Q^2 = -q^2 = -(E_f - E_i)^2 + \vec{q}^{\,2}$.

For the N$\Delta\beta$ vertices we define (see eq. (3.7)),

$$N^\Delta_\pi = \frac{ig_{\mathrm{N}\Delta\pi}F_{\mathrm{N}\Delta\pi}}{2m}\left[\frac{\mathcal{E}_\Delta\mathcal{E}}{(2\pi)^3 8\omega_\pi E_\Delta E}\right]^{\frac{1}{2}}, \qquad N^\Delta_\rho = \frac{g_{\mathrm{NN}\rho}G^{\mathrm{N}\Delta\rho}_{\mathrm{M}}}{2m}\frac{g_{\Delta\mathrm{N}\pi}}{g_{\mathrm{NN}\pi}}\left[\frac{\mathcal{E}_\Delta\mathcal{E}}{(2\pi)^3 8\omega_\rho E_\Delta E}\right]^{\frac{1}{2}} \tag{B5}$$

The interaction energies $\langle 0|b_\Delta(\vec{p}_) - \int d^3x\Gamma_{\Delta\mathrm{N}\beta}b^\dagger(\vec{p})|0\rangle$ for meson couplings to the N → Δ current are,

$$H_{\Delta\mathrm{N}\pi} = N^\Delta_\pi h_{\Delta\mathrm{N}\pi}\vec{\tau}_{\mathrm{N}\Delta}, \qquad H_{\Delta\mathrm{N}\rho} = N^\Delta_\rho h^\nu_{\Delta\mathrm{N}\rho}\epsilon_\nu\vec{\tau}_{\mathrm{N}\Delta} \tag{B6a}$$

$$h_{\Delta\mathrm{N}\pi} = -\vec{\sigma}_{\mathrm{N}\Delta}.\left[\vec{p}_\Delta\left(\frac{E}{m_\Delta} - \frac{(\vec{p}_\Delta.\vec{p})}{\mathcal{E}_\Delta m_\Delta}\right) - \vec{p}\right]\left[\mathbb{1} - (\frac{\vec{\sigma}.\vec{p}_\Delta}{\mathcal{E}_\Delta})(\frac{\vec{\sigma}.\vec{p}}{\mathcal{E}})\right] \tag{B6b}$$

$$\begin{aligned}h_{\Delta\mathrm{N}\rho} = {}& +\left[\frac{-\vec{p}_\Delta.\vec{\sigma}_{\mathrm{N}\Delta}}{m_\Delta};\vec{\sigma}_{\mathrm{N}\Delta} + \vec{p}_\Delta\frac{(\vec{\sigma}_{\mathrm{N}\Delta}.\vec{p}_\Delta)}{\mathcal{E}_\Delta m_\Delta}\right]^\mu \left[(\frac{\vec{\sigma}.\vec{p}_\Delta}{\mathcal{E}_\Delta}) + (\frac{\vec{\sigma}.\vec{p}}{\mathcal{E}});\vec{\sigma} + \frac{(\vec{\sigma}.\vec{p}_\Delta)\vec{\sigma}(\vec{\sigma}.\vec{p})}{\mathcal{E}_\Delta\mathcal{E}}\right]^\lambda \\ &\times\left[(p_\Delta - p)_\mu\delta^\nu_\lambda - (p_\Delta - p)_\lambda\delta^\nu_\mu)\right]\end{aligned} \tag{B6c}$$

The interaction energies $\langle 0|b(\vec{p}) - \int d^3x\Gamma_{\Delta\mathrm{N}\beta}b^\dagger_\Delta(\vec{p}_\Delta)|0\rangle$ for meson couplings to the Δ → N current are,

$$H_{\mathrm{N}\Delta\pi} = N^\Delta_\pi h_{\mathrm{N}\Delta\pi}\vec{\tau}^\dagger_{\mathrm{N}\Delta}, \qquad H_{\mathrm{N}\Delta\rho} = N^\Delta_\rho h^\nu_{\mathrm{N}\Delta\rho}\epsilon_\nu\vec{\tau}^\dagger_{\mathrm{N}\Delta} \tag{B7a}$$

$$h_{\mathrm{N}\Delta\pi} = +\left[\mathbb{1} - (\frac{\vec{\sigma}.\vec{p}_\Delta}{\mathcal{E}_\Delta})(\frac{\vec{\sigma}.\vec{p}}{\mathcal{E}})\right]\vec{\sigma}^\dagger_{\mathrm{N}\Delta}.\left[\vec{p}_\Delta\left(\frac{E}{m_\Delta} - \frac{\vec{p}_\Delta.\vec{p})}{\mathcal{E}_\Delta m_\Delta}\right) - \vec{p}\right] \tag{B7b}$$

$$\begin{aligned}h^\nu_{\mathrm{N}\Delta\rho} = {}& -\left[(\frac{\vec{\sigma}.\vec{p}_\Delta}{\mathcal{E}_\Delta}) + (\frac{\vec{\sigma}.\vec{p}}{\mathcal{E}});\vec{\sigma} + \frac{(\vec{\sigma}.\vec{p}_\Delta)\vec{\sigma}(\vec{\sigma}.\vec{p})}{\mathcal{E}_\Delta\mathcal{E}}\right]^\lambda \left[\frac{-\vec{\sigma}^\dagger_{\mathrm{N}\Delta}.\vec{p}_\Delta}{m_\Delta};\vec{\sigma}^\dagger_{\mathrm{N}\Delta} + \vec{p}_\Delta\frac{(\vec{\sigma}^\dagger_{\mathrm{N}\Delta}.\vec{p}_\Delta)}{\mathcal{E}_\Delta m_\Delta}\right]^\mu \\ &\times\left[(p - p_\Delta)_\mu\delta^\nu_\lambda - (p - p_\Delta)_\lambda\delta^\nu_\mu)\right]\end{aligned} \tag{B7c}$$

2. Currents for electromagnetic vertices

We define,

$$N_\gamma = \frac{e_p}{(2\pi)^3}\left\{\frac{\mathcal{E}_i\mathcal{E}_f}{4E_fE_i}\right\}^{1/2} \tag{B8}$$

The photon coupling to the positive frequency nucleon current $\langle 0|b(\vec{p}_f)\; J_{\mathrm{eff}}(0)b^\dagger(\vec{p}_i)|0\rangle$ is given by,

$$J^{\mu}_{\mathrm{NN}\gamma} = N_\gamma j^{\mu}_{\mathrm{NN}\gamma} \tag{B9a}$$

$$j^{0}_{\mathrm{NN}\gamma} = \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\Big[1 - \frac{E_f + E_i}{2m}\Big]\right] + \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\Big[1 + \frac{E_f + E_i}{2m}\Big]\right]\left[\frac{\vec{p}_f.\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i} + \frac{i\vec{\sigma}.(\vec{p}_f \times \vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i}\right] \tag{B9b}$$

$$\vec{j}_{\mathrm{NN}\gamma} = \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\right]\left[\frac{\vec{p}_f}{\mathcal{E}_f} + \frac{\vec{p}_i}{\mathcal{E}_i} + i\vec{\sigma} \times \left[\frac{\vec{p}_f}{\mathcal{E}_f} - \frac{\vec{p}_i}{\mathcal{E}_i}\right]\right] - \frac{\kappa F^{(2)}_{\mathrm{NN}\gamma}}{2m}(\vec{p}_f + \vec{p}_i)\left[1 - \frac{\vec{p}_f.\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i} - \frac{i\vec{\sigma}.(\vec{p}_f \times \vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i}\right] \tag{B9c}$$

The photon coupling to the pair-creation current $\langle 0|b(\vec{p}_f)\, J_{\mathrm{eff}}(0) d(-\vec{p}_i)|0\rangle$ is given by,

$$J_{\bar{\mathrm{N}}\mathrm{N}\gamma} = N_\gamma \epsilon_\mu j^{\mu}_{\bar{\mathrm{N}}\mathrm{N}\gamma} \tag{B10a}$$

$$j^{0}_{\bar{\mathrm{N}}\mathrm{N}\gamma} = \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\Big[1 + \frac{E_f - E_i}{2m}\Big]\right]\frac{\vec{\sigma}.\vec{p}_f}{\mathcal{E}_f} - \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\Big[1 - \frac{E_f - E_i}{2m}\Big]\right]\frac{\vec{\sigma}.\vec{p}_i}{\mathcal{E}_i} \tag{B10b}$$

$$\begin{aligned}\vec{j}_{\bar{\mathrm{N}}\mathrm{N}\gamma} = {} & \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\Big[1 - \frac{E_i}{m}\Big]\right]\vec{\sigma} + \frac{\kappa F^{(2)}_{\mathrm{NN}\gamma}}{2m}(\vec{p}_f + \vec{p}_i)\left[\frac{\vec{\sigma}.\vec{p}_f}{\mathcal{E}_f} + \frac{\vec{\sigma}.\vec{p}_i}{\mathcal{E}_i}\right] \\ & + \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\Big[1 + \frac{E_i}{m}\Big]\right]\left[\frac{i(\vec{p}_f \times \vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i} + \frac{\vec{\sigma}(\vec{p}_f.\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i} - \frac{\vec{p}_f(\vec{\sigma}.\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i} - \frac{(\vec{\sigma}.\vec{p}_f)\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i}\right]\end{aligned} \tag{B10c}$$

The photon coupling to the pair-annihilation current $\langle 0|d^\dagger(-\vec{p}_f)\, J_{\mathrm{eff}}(0) b^\dagger(\vec{p}_i)|0\rangle$ is given by,

$$J_{\mathrm{N}\bar{\mathrm{N}}\gamma} = N_\gamma \epsilon_\mu j^{\mu}_{\mathrm{N}\bar{\mathrm{N}}\gamma} \tag{B11a}$$

$$j^{0}_{\mathrm{N}\bar{\mathrm{N}}\gamma} = \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\Big[1 - \frac{E_f - E_i}{2m}\Big]\right]\frac{\vec{\sigma}.\vec{p}_i}{\mathcal{E}_i} - \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\Big[1 + \frac{E_f - E_i}{2m}\Big]\right]\frac{\vec{\sigma}.\vec{p}_f}{\mathcal{E}_f} \tag{B11b}$$

$$\begin{aligned}\vec{j}_{\mathrm{N}\bar{\mathrm{N}}\gamma} = {} & \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\Big[1 - \frac{E_f}{m}\Big]\right]\vec{\sigma} + \frac{\kappa F^{(2)}_{\mathrm{NN}\gamma}}{2m}(\vec{p}_f + \vec{p}_i)\left[\frac{\vec{\sigma}.\vec{p}_f}{\mathcal{E}_f} + \frac{\vec{\sigma}.\vec{p}_i}{\mathcal{E}_i}\right] \\ & + \left[F^{(1)}_{\mathrm{NN}\gamma} + \kappa F^{(2)}_{\mathrm{NN}\gamma}\Big[1 + \frac{E_f}{m}\Big]\right]\left[\frac{i(\vec{p}_f \times \vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i} + \frac{\vec{\sigma}(\vec{p}_f.\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i} - \frac{\vec{p}_f(\vec{\sigma}.\vec{p}_i)}{\mathcal{E}_f\mathcal{E}_i} - \frac{(\vec{\sigma}.\vec{p}_f)\vec{p}_i}{\mathcal{E}_f\mathcal{E}_i}\right]\end{aligned} \tag{B11c}$$

where the electromagnetic form form factors are defined as

$$F^{(1)}_{\mathrm{NN}\gamma} = \frac{1}{2}F^{(1);is}_{\mathrm{NN}\gamma} + \frac{1}{2}F^{(1);iv}_{\mathrm{NN}\gamma}\tau^0 \qquad \kappa_{\mathrm{N}}F^{(2)}_{\mathrm{NN}\gamma} = \frac{1}{2}\kappa^{is}F^{(2);iv}_{\mathrm{NN}\gamma} + \frac{1}{2}\kappa^{iv}F^{(2);iv}_{\mathrm{NN}\gamma}\tau^0 \tag{B12a}$$

$$\Rightarrow \text{protons}: \quad F^{(1)}_{\mathrm{NN}\gamma}(k^2 = 0) + \kappa_{\mathrm{N}}F^{(2)}_{\mathrm{NN}\gamma}(k^2 = 0) = (1) + (\frac{\kappa^{is} + \kappa^{iv}}{2}) = \mu_p \tag{B12b}$$

$$\Rightarrow \text{neutrons}: \quad F^{(1)}_{\mathrm{NN}\gamma}(k^2 = 0) + \kappa_{\mathrm{N}}F^{(2)}_{\mathrm{NN}\gamma}(k^2 = 0) = (0) + (\frac{\kappa^{is} - \kappa^{iv}}{2}) = \mu_n \tag{B12c}$$

For the N$\Delta\gamma$ vertices we define (see eq. (3.7)),

$$N^{\Delta}_{\gamma} = \frac{e_p}{(2\pi)^3 2m}\frac{G^{iv}_M}{2}\frac{g_{\Delta\mathrm{N}\pi}}{g_{\mathrm{NN}\pi}}\left[\frac{\mathcal{E}_\Delta\mathcal{E}}{4E_\Delta E}\right]^{\frac{1}{2}}; \qquad G^{iv}_M(0) = 1 + \kappa^{iv} = 4.706 \tag{B13}$$

The photon coupling to the N → Δ current $\langle 0|b_\Delta(\vec{p}_) J_{\mathrm{eff}}(0) b^\dagger(\vec{p})|0\rangle$ is given by,

$$J_{\Delta\mathrm{N}\gamma} = N^{\Delta}_{\gamma} j^{\nu}_{\Delta\mathrm{N}\gamma}\epsilon_\nu(\tau_{\mathrm{N}\Delta})^0 \tag{B14a}$$

$$\begin{aligned}j^{\nu}_{\Delta\mathrm{N}\gamma} = {} & + \left[\frac{-\vec{\sigma}_{\mathrm{N}\Delta}.\vec{p}_\Delta}{m_\Delta}; \vec{\sigma}_{\mathrm{N}\Delta} + \vec{p}_\Delta\frac{(\vec{\sigma}_{\mathrm{N}\Delta}.\vec{p}_\Delta)}{\mathcal{E}_\Delta m_\Delta}\right]^{\mu}\left[(\frac{\vec{\sigma}.\vec{p}_\Delta}{\mathcal{E}_\Delta}) + (\frac{\vec{\sigma}.\vec{p}}{\mathcal{E}}); \vec{\sigma} + \frac{(\vec{\sigma}.\vec{p}_\Delta)\vec{\sigma}(\vec{\sigma}.\vec{p})}{\mathcal{E}_\Delta\mathcal{E}}\right]^{\lambda} \\ & \times \left[(p_\Delta - p)_\mu\delta^\nu_\lambda - (p_\Delta - p)_\lambda\delta^\nu_\mu)\right]\end{aligned} \tag{B14b}$$

The photon coupling to the Δ → N current $\langle 0|b(\vec{p}) J_{\mathrm{eff}}(0) b^\dagger_\Delta(\vec{p}_\Delta)|0\rangle$ is given by,

$$J_{\mathrm{N}\Delta\gamma} = N^{\Delta}_{\gamma} j^{\nu}_{\mathrm{N}\Delta\gamma}\epsilon_\nu(\tau^\dagger_{\mathrm{N}\Delta})^0 \tag{B15a}$$

$$\begin{aligned}j^{\nu}_{\mathrm{N}\Delta\gamma} = {} & - \left[(\frac{\vec{\sigma}.\vec{p}_\Delta}{\mathcal{E}_\Delta}) + (\frac{\vec{\sigma}.\vec{p}}{\mathcal{E}}); \vec{\sigma} + \frac{(\vec{\sigma}.\vec{p}_\Delta)\vec{\sigma}(\vec{\sigma}.\vec{p})}{\mathcal{E}_\Delta\mathcal{E}}\right]^{\lambda}\left[\frac{-\vec{p}_\Delta.\vec{\sigma}^\dagger_{\mathrm{N}\Delta}}{m_\Delta}; \vec{\sigma}^\dagger_{\mathrm{N}\Delta} + \vec{p}_\Delta\frac{(\vec{\sigma}^\dagger_{\mathrm{N}\Delta}.\vec{p}_\Delta)}{\mathcal{E}_\Delta m_\Delta}\right]^{\mu} \\ & \times \left[(p - p_\Delta)_\mu\delta^\nu_\lambda - (p - p_\Delta)_\lambda\delta^\nu_\mu)\right]\end{aligned} \tag{B15b}$$

The photon coupling to the vector-meson-decay current is given by,

$$J^{\mu}_{\mathrm{VP}\gamma}(0)(q_{\mathrm{P}},q_{\mathrm{V}}) = \frac{-e_p g_{\mathrm{VP}\gamma}F_{\mathrm{VP}\gamma}}{m_{\mathrm{V}}}\frac{1}{(2\pi)^3}\frac{1}{\sqrt{4\omega_{\mathrm{V}}\omega_{\mathrm{P}}}}\epsilon^{\mu\xi ab}\epsilon_{\xi}(\vec{q}_{\mathrm{V}},\lambda_{\mathrm{V}})(q_{\mathrm{P}})_a(q_{\mathrm{V}})_b \tag{B16}$$

where $F_{\mathrm{VP}\gamma}(k^2=0)$=1, q_{V} and q_{P} are the 4-momenta delivered to the nucleons by the vector and pseudo-scalar mesons respectively.

APPENDIX C: THE MOMENTUM-SPACE NNγ RESCATTERING CALCULATION

The NNγ rescattering amplitudes in bremsstrahlung were first calculated by Brown [21] in r-space. For p-space calculations, a recipe has already been given in ref [10], although the residue terms are valid only for soft photons and the dominant final-state interaction is scaled by (guessed) off-shell minimal relativity factors and cast into the I-frame with a barycentric momentum k. This choice of frame does not minimize the effect of the neglected boost operators and the inclusion of minimal relativity factors disturbs the convergence properties of the p-space integral.

Casting the entire rescattering amplitude into the A-frame of eq (3.33) requires a somewhat different numerical procedure. We start from eq. (3.44) and simplify our notation by defining,

$$\mathcal{M}^{\mathrm{R}}_{fi} = K\int\!\!\int H(\hat{p})d\theta d\phi \qquad\qquad H(\hat{p}) = \int\frac{F(\vec{p})}{G_i(\vec{p})G_f(\vec{p})}dp \tag{C1}$$

with,

$$\begin{aligned}
F(\vec{p}) &= p^2\sin\theta[E(\vec{p}_3+\frac{1}{4}\vec{k})+E(\vec{p}+\frac{1}{4}\vec{k})][E(\vec{p}_1-\frac{1}{4}\vec{k})+E(\vec{p}-\frac{1}{4}\vec{k})]\\
&\quad\times\sum_{M_S M'_S}\langle\vec{p}_3\,\widetilde{+}\,\frac{1}{4}\vec{k};S_f M_{S_f};T_f M_T|[1+i\chi(-\frac{\vec{k}}{2})]t^{(-)\dagger}[1-i\chi(-\frac{\vec{k}}{2})]|\vec{p}+\frac{1}{4}\vec{k};S_f M'_S;T_f M_T\rangle\\
&\quad\times\langle S_f M'_S;T_f M_T|J_{\mathrm{NN}\gamma}[1](-\vec{p}+\frac{\vec{k}}{2},-\vec{p}-\frac{\vec{k}}{2})|S_i M_S;T_i M_T\rangle\\
&\quad\times\langle+\vec{p}-\frac{1}{4}\vec{k};S_i M_S;T_i M_T|[1+i\chi(+\frac{\vec{k}}{2})]G_i t^{(+)}[1-i\chi(+\frac{\vec{k}}{2})]|\vec{p}_1\,\widetilde{-}\,\frac{1}{4}\vec{k};S_i M_{S_i};T_i M_T\rangle\\
G_i(\vec{p}) &= (\vec{p}_1-\frac{1}{4}\vec{k})^2-(\vec{p}-\frac{1}{4}\vec{k})^2+i\eta_i = \Delta_i^2-(p-\frac{1}{4}|\vec{k}|\cos\vartheta)^2+i\eta_i\\
G_f(\vec{p}) &= (\vec{p}_3+\frac{1}{4}\vec{k})^2-(\vec{p}+\frac{1}{4}\vec{k})^2+i\eta_f = \Delta_f^2-(p+\frac{1}{4}|\vec{k}|\cos\vartheta)^2+i\eta_f\\
\Delta_i^2 &= \left(\frac{1}{4}|\vec{k}|\cos\vartheta\right)^2-\vec{p}_1.\vec{p}_2 \qquad\qquad \Delta_f^2 = \left(\frac{1}{4}|\vec{k}|\cos\vartheta\right)^2-\vec{p}_3.\vec{p}_4\\
K &= \frac{N}{2}(-1)^{(S_f+S_i+T_f+T_i)}
\end{aligned} \tag{C2}$$

with $\cos\vartheta=\hat{p}.\hat{k}=\sin\theta_p\sin\theta_k\cos(\phi_p-\phi_k)+\cos\theta_p\cos\theta_k$. Within the (maximally symmetric) A-frame the effects of the boost operators $\chi(+\frac{1}{2}\vec{k})$ and $\chi(-\frac{1}{2}\vec{k})$ can be expected to substantially cancel and will therefore be neglected in the present numerical applications. Their inclusion would change nothing that we will discuss in this appendix.

For $\Delta_i,\Delta_f<0$, there are no poles on the real axis, but it is easy to see that there are singularities in the Greens' functions at $\vec{p}=\vec{p}_1$, $\vec{p}_2$, $\vec{p}_3$, $\vec{p}_4$ as well as $(\vec{p}_1-\vec{p}).(\vec{p}_2-\vec{p})$ and $(\vec{p}_3-\vec{p}).(\vec{p}_4-\vec{p})$, the latter two requiring particular attention when $\cos\vartheta_i=\pm4\sqrt{\vec{p}_1.\vec{p}_2}/|\vec{k}|$ or $\cos\vartheta_f=\pm4\sqrt{\vec{p}_3.\vec{p}_4}/|\vec{k}|$ cause $\Delta_i=0$ or $\Delta_f=0$, so that G_i^{-1} and/or G_f^{-1} each have a 2^{nd}-order pole (on the real axis). We adopt the utilatarian attitude of noting that for experiments below the π-production threshold, the second-order poles only occur at $\theta_1^{\mathrm{Lab}},\theta_2^{\mathrm{Lab}}<6°$, and even then, only for $\theta_\gamma^{\mathrm{Lab}}<1°$. Since no data exists in this region, we simply defer a treatment of second order singularities and confine our attention to the kinematics containing the simple and separable poles,

$$\begin{aligned}
p &= p_i^{(\pm)} = +\frac{1}{4}|\vec{k}|\cos\vartheta\pm\Delta_i \qquad\qquad \text{if}\ \ \Delta_i^2\geq0\ \ \text{then}\ \ \Delta_i=|p_i^{(\pm)}-\frac{1}{4}|\vec{k}|\cos\vartheta|\\
p &= p_f^{(\pm)} = -\frac{1}{4}|\vec{k}|\cos\vartheta\pm\Delta_f \qquad\qquad \text{if}\ \ \Delta_f^2\geq0\ \ \text{then}\ \ \Delta_f=|p_f^{(\pm)}+\frac{1}{4}|\vec{k}|\cos\vartheta|
\end{aligned} \tag{C3}$$

and evaluate Δ_i^2 and Δ_f^2 to determine if poles exist on the positive real $p-$axis. Defining such poles to be vectors so that $\hat{p}_i^{\pm}{=}\hat{p}_f^{\pm}{=}\hat{p}$, we obtain,

$$H(\hat{p}) = \text{V.P.} \int dp \frac{F(\vec{p})}{\left[\Delta_i^2 - (p - \frac{1}{4}|\vec{k}|\cos\vartheta)^2\right]\left[\Delta_f^2 - (p + \frac{1}{4}|\vec{k}|\cos\vartheta)^2\right]} - Z \tag{C4}$$

where

$$Z = \frac{i\pi F(\vec{p}_i^{\,(+)})}{2\Delta_i G_f(p_i^{(+)})} + \frac{i\pi F(\vec{p}_i^{\,(-)})}{2\Delta_i G_f(p_i^{(-)})} + \frac{i\pi F(\vec{p}_f^{\,(+)})}{2\Delta_f G_i(p_f^{(+)})} + \frac{i\pi F(\vec{p}_f^{\,(-)})}{2\Delta_f G_i(p_f^{(-)})} \tag{C5}$$

remains well defined. This formally completes the specification of the integral. However, for practical purposes, we need to add special forms of zero to smooth the divergences near the poles. With a simple generalization of

$$\text{V.P.} \int_{a_1}^{a_2} \frac{1}{k^2 - p^2} dp = \frac{1}{2k} \ln|\frac{a_2 + k}{a_2 - k} \cdot \frac{k - a_1}{k + a_1}| \tag{C6}$$

we obtain,

$$H = H_{p_i^{(+)}} + H_{p_i^{(-)}} + H_{p_f^{(+)}} + H_{p_f^{(-)}} \tag{C7}$$

where

$$H_{p_i^{(+)}} = \int_{a1}^{a2} \frac{1}{G_i(\vec{p})} \left[\frac{F(\vec{p})}{G_f(\vec{p})} - \frac{F(\vec{p}_i^{\,(+)})}{G_f(\vec{p}_i^{\,(+)})}\right] dp + \frac{1}{2\Delta_i} \frac{F(\vec{p}_i^{\,(+)})}{G_f(p_i^{(+)})} \left[\ln|\frac{(a2 - p_i^{(-)})}{(a2 - p_i^{(+)})} \frac{(a1 - p_i^{(+)})}{(a1 - p_i^{(-)})}| - i\pi\right] \tag{C8a}$$

$$H_{p_i^{(-)}} = \int_{b1}^{b2} \frac{1}{G_i(\vec{p})} \left[\frac{F(\vec{p})}{G_f(\vec{p})} - \frac{F(\vec{p}_i^{\,(-)})}{G_f(p_i^{(-)})}\right] dp + \frac{1}{2\Delta_i} \frac{F(\vec{p}_i^{\,(-)})}{G_f(p_i^{(-)})} \left[\ln|\frac{(b2 - p_i^{(-)})}{(b2 - p_i^{(+)})} \frac{(b1 - p_i^{(+)})}{(b1 - p_i^{(-)})}| - i\pi\right] \tag{C8b}$$

$$H_{p_f^{(+)}} = \int_{c1}^{c2} \frac{1}{G_f(\vec{p})} \left[\frac{F(\vec{p})}{G_i(\vec{p})} - \frac{F(\vec{p}_f^{\,(+)})}{G_i(\vec{p}_f^{\,(+)})}\right] dp + \frac{1}{2\Delta_f} \frac{F(\vec{p}_f^{\,(+)})}{G_i(p_f^{(+)})} \left[\ln|\frac{(c2 - p_f^{(-)})}{(c2 - p_f^{(+)})} \frac{(c1 - p_f^{(+)})}{(c1 - p_f^{(-)})}| - i\pi\right] \tag{C8c}$$

$$H_{p_f^{(-)}} = \int_{d1}^{d2} \frac{1}{G_f(\vec{p})} \left[\frac{F(\vec{p})}{G_i(\vec{p})} - \frac{F(\vec{p}_f^{\,(-)})}{G_i(p_f^{(-)})}\right] dp + \frac{1}{2\Delta_f} \frac{F(\vec{p}_f^{\,(-)})}{G_i(p_f^{(-)})} \left[\ln|\frac{(d2 - p_f^{(-)})}{(d2 - p_f^{(+)})} \frac{(d1 - p_f^{(+)})}{(d1 - p_f^{(-)})}| - i\pi\right] \tag{C8d}$$

where the domains [a1:a2] + [b1:b2] + [c1:c2] + [d1:d2] span [0:∞] and contain the poles $p_i^{(+)}$, $p_i^{(-)}$, $p_f^{(+)}$ and $p_f^{(-)}$ respectively. In the event that $p_\alpha^{(\pm)}$ does not exist in [0:∞], then our expressions require $F(p_\alpha^{(\pm)}) = 0$.

---

[1] J. A. Eden and M. F. Gari, Phys. Lett. **B347**, 187 (1995);

[2] V. Herrmann and K. Nakayama, O. Scholten, and H. Arellano, Nucl. Phys. **A582**, 568 (1995).

[3] M. Gari and H. Hyuga, Z. Phys **A277**, 291, (1976); Nucl. Phys. **A278**, 372 (1977).

[4] F. E. Low, Phys. Rev. **110**, 974 (1958).

[5] S. L. Adler and Y. Dothan, Phys. Rev. **151**, 1267 (1966).

[6] E. M. Nyman, Phys. Rev. **170**, 1628; Phys Lett **25B** (1967) 135 (1968).

[7] G. Felsner, Phys. Lett. **25B**, 290 (1967).
[8] R. L. Workman and H. W. Fearing, Phys. Rev. **C34**, 780 (1986).
[9] V. R. Brown, P. L. Anthony, and J. Franklin, Phys. Rev. **C44**, 1296 (1991).
[10] V. Herrmann, J. Speth, and K. Nakayama, Phys. Rev. **C43**, 394 (1991).
[11] V. Herrmann, and K. Nakayama, Phys. Rev. **C46**, 2199 (1992).
[12] H. W. Fearing, TRIUMF Preprint TRI-PP-93-12 (March 1993).
[13] H. W. Fearing, in Kernfysisch Versneller Instituut Report KVI-180i, 1992, (unpublished).
[14] J. A. Eden, D. Plümper, M. F. Gari and H. Hebach, Z. Phys. **A347** (1993) 145.
[15] J. A. Eden and M. F. Gari, Ruhr Universität Bochum Report RUB-MEP-81/94; nucl-th/9501034.
[16] A. Katsogiannis and K. Amos, Phys. Rev. **C47**, 1376 (1993).
[17] M. Jetter, H. Freitag, and H. von Geramb, Phys. Scr. **48**, 229 (1993).
[18] M. Jetter and H.V. von Geramb, Phys. Rev.**C49**, 1832 (1994).
[19] M. Jetter and H. Fearing, TRIUMF preprint TRI-PP-94-90; nucl-th/9410040.
[20] F. de. Jong, K. Nakayama, V. Herrmann and O. Scholten, Phys. Lett **B333**, 1 (1994); F. de. Jong, K. Nakayama and T. -S. H. Lee, nucl-th/9412013.
[21] V. R. Brown, Phys. Rev. **177**, 1498 (1969).
[22] V. R. Brown, Phys. Lett. **B23**, 259 (1971).
[23] J. Flender and M. F. Gari, Phys. Rev. **C51**, R1 (1995); Z. Phys. **A343**, 467 (1992).
[24] T. D. Cohen, Phys. Rec **D34**, 2187 (1986); N. Kaiser, U. G. Meissner and W. Weise, Phys. Lett. **198B**, 75 (1988).
[25] P. Alberto, E. Ruiz Arriola, M. Fiolhais, F. Grümmer, J. N. Urbano and K. Goeke, Phys. Lett. **208B**, 75 (1988).
[26] B. L. G. Bakker, B. Brozoian, J. N. Maslow and H. J. Weber, Phys. Rev **C25**, 1134 (1982).
[27] M. F. Gari and W. Krümpelmann, Phys. Rev. **D45**, 1817 (1992).
[28] D. Plümper, J. Flender and M. F. Gari, Phys. Rev. **C49**, 2370 (1994).
[29] D. Huber, H.Witala, H. Kamada and W. Glöckle, Proceedings of XVth European conference on Few-Body Problems in Physics, FBXV Pensicola, Spain (1995).
[30] A. W. Thomas and K. Holinde, Phys. Rev. Lett **63**, 2025 (1989).
[31] M. Gari, H. Hyuga and B. Sommer, Phys. Rev. **C14**, 2169, (1976);
[32] G. -H. Niephaus, M. Gari and B. Sommer, Phys. Rev. **C20**, 1096, (1979);
[33] M. Gari, G. -H. Niephaus and B. Sommer, Phys. Rev. **C23**, 504, (1981);
[34] W. N. Cottingham, M. Lacombe, B. Loiseau, J. M. Richard and R. Vinh Mau, Phys. Rev. **D8**, 800 (1973); M. Lacombe, B. Loiseau, J. M. Richard, R. Vinh Mau, J. Côté, P. Pirés and R. de Tourreil, Phys. Rev. **C21**, 861 (1980).
[35] K. Michaelian et al, Phys. Rev. **C41**, 2689 (1990). Phys. Rev. **C21**, 861 (1980).
[36] D. Drechsel and L. V. Maximon, Annals of Phys. **49** 403 (1968).
[37] D. O. Riska, "Exchange Currents", Preprint Series in Theoretical Physics, HU-TFT-89-15, University of Helsinki
[38] M. Araki and N. Kamal, Phys. Rev. **D29**, 1345 (1984).
[39] C. D. de Tar,Phys. Rev. **D24**, 752 (1981); Phys. Rev. **D24**, 762 (1981).
[40] S. Deister, M. F. Gari, W. Krümpelmann and M. Mahlke, Few Body Syst. **10**,1 (1991).
[41] U. Karlfuß and M. F. Gari, Nucl Phys **A408**, 507 (1983).
[42] M. Chemtob, E. Moniz, and M. Rho, Phys. Rev. **C10**, 344 (1974).
[43] E. Hummel and J. A. Tjon, Phys. Rev. Lett. **63**,1788 (1989).
[44] Particle Data Group, Rev. Mod. Phys. **52**,2II, 81 (1980); O. Dumbrajs, R. Koch, H. Pilkuhn, G. C. Oades, H. Behrens, J. J. de Swart and P. Knoll, Nucl. Phys. **B216**, 277 (1983).
[45] G.E.Brown and W.Weise, Phys. Rep. **22C**, 279 (1975).
[46] A.M.Green *et al*, Rep. Prog. Phys **39** 1109 (1976)
[47] T.Ericson and W.Weise, *Pions and Nuclei*, Clarendon Press, Oxford (1988).
[48] H. Arenhövel, Nucl. Phys. **A247**, 473 (1975).
[49] R. Machleidt, K. Holinde and C. Elster, Phys. Rep. **149**, 1 (1987); R. Machleidt, Adv. Nucl. Phys **19**, 189 (1989).
[50] L. L. Foldy, Phys. Rev. **122**, 275 (1961).
[51] R. A. Krajcik and L. L. Foldy, Phys. Rev. Lett. **24**, 545 (1970); Phys. Rev. **D10**, 1777 (1974).
[52] J. L. Friar, Phys. Rev. **C12**, 695 (1975). Phys. Rev. **D10**, 1777 (1974).
[53] W. Glöckle and L. Müller, Phys. Rev. **C23**, 1183 (1980).
[54] M. M. Nagels T. A. Rijken and J. J. de Swart, Phys. Rev. **D17**, 768 (1978).
[55] S. Weinberg, Phys. Rev. Lett. **18**, 188 (1967).
[56] J. Wess and B. Zumino, Phys. Rev. **163**, 1727 (1967).
[57] F. Gross, Phys. Rev. **C26**, 2203 (1982).
[58] F. Gross, K. M. Maung, J. A. Tjon, L. W. Townsend and S. J. Wallace, Phys. Rev. **C40**, R10 (1989).
[59] F. Gross, J. W. Van Orden and K. Holinde, Phys. Rev. **C41**, R1909 (1990).
[60] R. M. Woloshyn and A. D., Jackson, Nucl. Phys. **B64** 269 (1973).
[61] J. Fleischer and J. A. Tjon, Nucl. Phys. **B84**, 375 (1975).
[62] E. Erkelenz, Phys. Rep. **C13** 193 (1974).